OPEN ACCESS

Giant Fibrovascular Vulval Polyp in a Young Female: A Case Report



Rena Helmi¹, Wassan Nori² and Zeena Helmi^{2,*}

¹Department of Dermatology, College of Medicine, Mustansiriyah University, Baghdad, Iraq ²Department of Gynecology and Obstetrics, College of Medicine, Mustansiriyah University, Baghdad, Iraq

Abstract:

CASE REPORT

Background: Fibrovascular stromal polyps (fibroepithelial polyps; FEPs) are benign tumors occurring in many parts of the body and are infrequently reported in the vulva. They tend to be small, but few present in giant sizes, which adds diagnostic challenges.

Case presentation: A 24-year-old female with a five-year history of a right vulval mass that reached a massive size $(35 \times 15 \times 10 \text{ cm})$, complained of a large, painless, soft, pedunculated mass causing discomfort and interference with daily activities. On examination, the mass was firm, not tender, and associated with surface ulceration and infection. The mass was surgically removed and subjected to histological examination to exclude any malignancy. The observation indicated the presence of a fibroepithelial (fibrovascular) polyp; thus, follow-up was advised.

Conclusion: This case report has explored giant FEPs' presentation, diagnosis, therapeutic approaches, and possible consequences, and comprehensively discussed the findings of earlier studies. A better understanding of giant FEPs can improve their management and guide prescription for tailored treatment, which must involve evaluation of the psychological impact on patient mental health for a comprehensive holistic patient approach.

Keywords: Fibroepithelial polyp, Giant, Benign vulva neoplasm, Diagnosis, Complication, Tumors.

© 2025 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: https://creativecommons.org/licenses/by/4.0/legalcode. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

* Address correspondence to this author at the Department of Gynecology and Obstetrics, College of Medicine, Mustansiriyah University, Baghdad, Iraq; Tel: 009647713726331; E-mail: zeena.helmi@uomustansiriyah.edu.iq

Cite as: Helmi R, Nori W, Helmi Z. Giant Fibrovascular Vulval Polyp in a Young Female: A Case Report. Open Dermatol J, 2025; 19: e18743722377696. http://dx.doi.org/10.2174/0118743722377696250324062719



Received: December 15, 2024 Revised: March 05, 2025 Accepted: March 10, 2025 Published: April 10, 2025



Send Orders for Reprints to reprints@benthamscience.net

1. INTRODUCTION

Fibrovascular stromal polyps (fibroepithelial polyps; FEPs), which are non-malignant in nature, commonly occur in various anatomical locations. They are typically located in the skin folds, such as the neck, armpit, breast, groin, oral cavity, bladder neck, and genital regions [1-9]. They impact approximately 25% of the population, and their prevalence rises with age [10].

FEPs exhibit a range of clinical manifestations, ranging from minor to big pedunculated tumors. FEPs are typically small in size. They are referred to as "giants" when they exceed 5 cm in any dimension [11]. However, some literature studies use a cut-off value of 10 cm to define giant FEP [12].

Vulvar occurrences of FEPs are uncommon, and the manifestation of giant fibroepithelial polyps (GFEPs) in this area is exceptionally unusual, as documented in the literature. As they are very big and affect nearby tissues, GFEPs cause noticeable symptoms, like vulvar distention and pain and problems with daily activities, thus limiting intimate functions [13].

Ulceration, hemorrhaging, and secondary infections have also been reported. In exceptionally large cases,

lymphedema occurs due to impaired lymphatic drainage. The physical pain caused by GFEPs can have a detrimental effect on patient's body perception, resulting in feelings of anxiety and depression, thus imposing a psychological impact on their lives [14, 15].

We, herein, present the case of a young female who presented with a vulval mass for five years, and embarrassment and social restrictions have refrained her from early counseling.

2. CASE REPORT

A 24-year-old woman who was sexually inactive visited the clinic with a complaint of a large, painless, soft, pedunculated mass. The mass extended from the right labia majora to below the middle of the thigh and caused discomfort during daily activities. The mass progressively enlarged over a span of five years. There was a negative history of vaginal discharge, fever, weight loss, or issues related to the bowel or bladder.

The patient experienced menarche at the age of 12 and had regular menstrual cycles with a normal flow. Her previous medical and surgical records were unremarkable. The patient was a virgin with no prior history of sexually transmitted diseases, genital injuries, drug use, smoking, or alcohol consumption.



Fig. (1). Local examination revealed a $35 \times 15 \times 10$ cm soft, pedunculated mass extending from the right labia majora causing labial elongation and extending below the midthigh.

The patient had concerns that medical intervention might compromise her virginity. Due to religious and cultural restrictions and the associated taboos surrounding virginity, she refrained from seeking any medical guidance until the condition had grown to an intolerable size.

Upon local inspection, a soft, pedunculated mass of 35 \times 15 \times 10 cm was seen. The mass originated from the

right labia majora, resulting in an extension of the labia and further extending to the midthigh. In an upright position, the mass was located a few millimeters above the patella (Fig. 1). The mass was warm to the touch, not tender, and had a layer of skin with two ulcers on top, measuring 2×1 cm and 1×1 cm, respectively, with a yellowish discharge. The mons and the labia on the opposite side appeared normal. The trans-illumination test yielded a negative result. There was no swelling of the lymph nodes and no connection with the inguinal canal. The general inspection yielded no notable findings. The hematological and biochemical studies yielded normal The ultrasound examination revealed no results. abnormalities in the pelvic or uterine regions, so the decision was made to proceed with immediate surgical removal of the tumor without a preoperative MRI or biopsy.



Fig. (2). The mass was excised completely from its base and then ligated using Vicryl number 0 suture.



Fig. (3). Resected perineal mass weighing 850 grams.

Authors, Year\Refs	Age	Site	Size/cm	Duration	Presenting Symptoms
The current case, 2024	24	Rt labia majora	35×15×10	5 years	Ulcerating pedunculated mass
Korkontzelos et al., 2023 [16]	45	Rt labia majora	10×13	2 years	Pedunculated, painless mass that was inflamed and ulcerated
Dura <i>et al.,</i> 2023 [17]	21	Rt labia majora	3×6×9	2 years	Painless mass causing discomfort and refrainment from sex
Martínez, 2023 [18]	30	Rt labia majora	20×15	1 year	Pedunculated mass with pulling pain
Martínez, 2023 [18]	36	Lt labia majora	7×5×5	1 year	A pedunculated mass associated with a pressure ulcer
Mulik and Acharya, 2023 [19]	56	Rt labia majora	8×7.5×7.5	5 years	A pedunculated mass interfering with walking
Donata, 2022 [20]	21	Lt labia majora	12×4	3	A painless mass causing discomfort and refrainment from sex
Kurniawati, 2022 [21]	23	Rt labia majora	16×11×6	1.6 years	A painless mass causing discomfort
Kurniadi <i>et al.,</i> 2022 [12]	28	Lt and Rt labia majora	Lt 20×18×8 Rt 9×4×2	4 years	A painless firm mass with leukorrhea
Smet <i>et al.</i> [22]	22	Rt vulva	15×11×4	11 years	A mass involving bleeding from giant pedunculated multiple finger- like projections.
Quinn et al., 2020 [23]	24	Rt labia majora	5×6	2 years	Gelatinous polyp with ulceration and bleeding
Yoo J et al., 2019 [24]	14	Rt labia majora	20×12.5× 3.5	3 years	A huge, exophytic, polypoidal mass raised from Rt labium majora
Rexhepi et al., 2018 [13]	16	Rt labia majora	18×12×3	2 years	A giant polypoid mass interfering with daily activities
Kumar et al., 2017 [25]	20	Lt labia majora	42×22×10	10 years	A giant mass preventing walking and interfering with urination
Chawla et al., 2017 [11]	32	Lt labia majora	10×10	10 years	A pedunculated polypoidal painful mass associated with ulcer
Colak Elif, 2015 [26]	43	Lt perineum	18×12	29 years	A growing mass with discomfort
Madueke, 2013 [27]	21	Rt labia majora	18.5	6 months	A grapefruit-sized ulcerating pedunculated mass

Table 1. Summary of published FEPs cases in the last decade.

Local anesthesia was administered to the patient because she refused general or spinal anesthesia. After admission to the hospital, the patient was lithotomy positioned. The procedure was performed under aseptic conditions. The resection was carried out surgically, followed by ligation of the wide pedicle (2.5 cm). Adequate hemostasis was achieved (Fig. 2). The patient was prescribed amoxicillin/clavulanic acid at a dosage of 1 gram orally twice daily and was discharged after a few hours. After a period of two weeks, the patient came for a follow-up visit; the examination revealed a fully healed wound without any complications. The patient expressed satisfaction with the outcome. There was no sign of recurrence.



Fig. (4). Microphotograph showing benign-looking squamous epithelium (red arrow) with underlying fibrovascular stroma (green arrow); it was consistent with fibrovascular polyp (angiomatous). H&E, X100.

Macroscopically, the removed mass was a solitary, homogenous mass with a smooth skin surface, having dimensions of $35 \times 15 \times 10$ cm and weighing 850 g, as shown in Fig. (3). The sample was subjected to histopathological analysis.

Under microscopic examination, the sections revealed benign squamous epithelium tissue covering the cystic wall, made of fibrous tissue with blood-filled blood vessels of various sizes. These vessels were lined by benign endothelial cells and accompanied by chronic inflammatory cell infiltration in certain areas. The observation indicated the presence of a fibroepithelial (fibrovascular) polyp, as depicted in Figs. (4 and 5). No cancerous cells or abnormal cells were detected.



Fig. (5). Microphotograph showing benign-looking squamous epithelium (red arrow) with underlying fibrovascular stroma (green arrow); it was consistent with fibrovascular polyp (angiomatous). H&E, X200.

3. DISCUSSION

A giant fibroepithelial polyp is an uncommon vulvar tumor that arises from the mesenchyme. Although it is noncancerous, it mimics a malignant tumor. GFEPs reported in the literature in the last decade are summarized in Table 1.

In the earlier works, these GFEP tumours have been predominantly seen in the labia majora, vulva, and perineum. This can be attributed to hormonal sensitivity in these areas. The subepithelial stroma in the mentioned areas exhibits sensitivity to estrogen and progesterone, promoting the development and growth of these polyps. Many are indeed seen among reproductive-age women, further supporting this theory.

Another probable cause is that the vulvar area experiences friction and irritation. These mechanical stimulations may trigger the formation and expansion of these polyps in these anatomical sites. Characteristically, FEPs have a pedunculated nature, rubbery consistency, and often ulcerated surfaces, supporting the diagnosis. Vulvoscopy and biopsy play a crucial role in differentiating GFEPs from other vulvar lesions, particularly those of malignant potential [28]. Moreover, endoscopy has been reported to be useful in differentiating FEPs originating in the bladder and manifesting as hematuria [29].

The diagnosis of GFEPs is performed by histopathological examination of tissue samples, which is the gold standard for confirming the condition and ruling out other possibilities [30].

An accurate diagnosis usually requires further investigation; imaging modalities, such as CT or MRI, are sometimes employed for GFEPs to determine where the tumor originated and where it extends by looking at its blood supply and circulation, preferably pre-operatively. In addition, they help exclude other differential diagnoses, particularly those of malignant nature. Kato *et al.* highlighted that the identification of adipose tissue on MRI is a distinct observation that indicates FEPs [31]. Additionally, Yoo *et al.* declared observing adipose cell accumulation in close proximity to the stalk of FEPs, indicating MRI as beneficial in identifying fat [24]. Fig. (6) summarizes the diagnostic steps involved in detecting FEPs.

Some of the differential diagnoses associated with GFEPs of the vulva are presented as follows:

3.1. Condylomata Acuminata

It tends to be smaller in size, has usually multiple lesions, and indicates positive HPV testing.

3.2. Angiomyxoma

It tends to have a deeper location and a closer coloration.

3.3. Vulvar Sarcoma

It tends to grow faster with an irregular border and deeper involvement.

Other less common diagnoses include squamous cell carcinoma, embryonal rhabdomyosarcoma, granular cell tumors, neurofibromas, juvenile xanthogranuloma, and leiomyomas [14, 32, 33].

At the moment, the predisposing factors for GFEPs are not entirely understood. Unraveling the pathogenesis and etiology is critical for a comprehensive understanding of GFEPs. Certain hypotheses posit hormonal influences. Estrogen and progesterone fluctuations during pregnancy, hormone replacement therapy, and the use of combined birth control pills have been recognized to stimulate the growth of GFEPs' epithelium, which has been proven to be responsive to hormones. Moreover, there are reports of tumor size reduction after pregnancy. Hormonal theory might explain why the majority of cases occur during the reproductive years [12, 24, 34-36].



Fig. (6). Flow chart for diagnosing FEPs.

Some scholars have linked hyperlipidemia, insulin resistance, and obesity as metabolic disorders with GFEPs [37].

Additionally, FEPs have been associated with people suffering from chronic inflammatory conditions, including Crohn's disease, psoriasis, and inherited lymphedema. Preliminary research indicates that their development may have a genetic component. Still, further research is needed [38-40].

The main approach to the treatment of GFEPs is surgical excision. Methods vary from basic excision to sequential excision for bigger polyps. It may be appropriate to pursue minimally invasive methods, such as laser ablation or electrocautery, for smaller lesions [16].

Surgical procedures involving extensive tissue removal may result in delayed wound healing, the formation of scars, and susceptibility to infections. Lymphedema manifestation entails the continuous utilization of compression garments and lymphedema treatment [21], [41]. A meticulous implementation of surgical methods, perioperative treatment, and the provision of suitable postoperative care for lymphedema are supposed to reduce these risks.

It is worth mentioning that GFEPs in pregnancy may impose therapeutic challenges, as they tend to grow rapidly and may bleed. Smet *et al.* [22] reported a case that presented in term pregnancy with a bleeding attack that called for an urgent surgical intervention to remove GFEPs of $15 \times 11 \times 4$ cm and secure hemostasis. In order not to compromise the site of surgery, the decision was to deliver the baby *via* C-section.

It is important to remember that GFEPs can adversely impact the quality of life of the individual. The patient endures prolonged pain with the possibility of scarring and recurrence after the surgery, which can also have a negative effect on general well-being and sexual health. The presence of anxiety and depression linked to the initial diagnosis and its sequelae might persist even after the surgery. For that reason, it is crucial to pursue effective therapies for psychological components and provide emotional assistance to the patient [27, 42-45].

Although surgical excision is usually effective in treating GFEPs, a small number of cases may experience a recurrence, especially in cases with incomplete resection. Malignant transformation, although extremely rare, should still be considered. Regular post-treatment check-ups are essential for evaluating the possibility of recurrence and resolving any potential problems or malignant changes [13].

The heterogeneity of GFEP sizes and locations makes their identification challenging. It is still not clear why, in some cases, hormones are blamed for fast polyp growth, as in adolescence and pregnancy, while in other cases, the mass grows slowly for a while and then grows quickly for no clear reason [12]. To the best of our knowledge, the largest vulval GFEP reported has been of 42 cm [7]. The current case is the second-largest size ever reported in the literature.

Most of the published studies have been retrospective and case reports, including small numbers, making it difficult to draw definitive conclusions. Investigating GFEPs' pathogenesis, prognosis, long-term outcomes, and quality-of-life considerations can help develop a thorough understanding of these fascinating tumors. Continued study is necessary to improve management techniques and deepen our comprehension of these distinct vulvar tumors.

CONCLUSION

Giant fibrovascular vulvar polyps in young females are rare, but significant benign growths that require careful clinical evaluation and management. These polyps typically present as large, often painless masses in the vulvar region and can be associated with mechanical stress or hormonal influences. Early diagnosis is crucial for appropriate intervention, which may involve surgical excision to prevent complications, such as discomfort, ulceration, or potential malignancy. While most cases are benign, understanding their pathophysiology and considering differential diagnoses, such as tumors or infections, are important in establishing the correct treatment plan. Further research is needed to explore the underlying causes, risk factors, and long-term outcomes of these uncommon yet impactful lesions in young women.

AUTHORS' CONTRIBUTIONS

All authors have contributed equally to the manuscript and agreed to the submission of the final version.

LIST OF ABBREVIATIONS

СТ	=	Compound Tomography						
HPV	=	Human Papilloma Virus						
MRI	=	Magnatic Resonance Imaging						
GEPs	=	Giant Fibroepitheial Polyps						
Cm	=	Centimeter						
FEPs	=	Fibroepitheial Polyps						

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study obtained ethical approval from the ethics committee of Mustansiriyah University, Baghdad, Iraq (ref. no. MOG 37; dated 6/2024).

HUMAN AND ANIMAL RIGHTS

All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

An informed consent was obtained from the patient before performing the study.

STANDARDS OF REPORTING

CARE guidelines were followed.

AVAILABILITY OF DATA

The authors confirm that the data supporting the findings of this research are available within the article.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

The authors would like to thank the College of Medicine, Al-Musnasiriyah University, Iraq.

REFERENCES

- [1] Navada MH, Bhat PRB, Rao SV, G N. Large fibroepithelial polyp of vulva. Case Rep Dermatol Med 2011; 2011: 1-2. http://dx.doi.org/10.1155/2011/273181 PMID: 23213577
- [2] Arora BK. A clinical case of large fibroepithelial polyp of breast nipple. Int J Case Rep Images 2019; 10: 100997Z01BA2019.
- [3] Raphael Avidime A, Usman H. Bilateral giant fibroepithelial labial mass: A case report. J Obstet Gynaecol Can 2017; 39(7): 564-6. http://dx.doi.org/10.1016/j.jogc.2017.02.008 PMID: 28483334
- [4] Leffelman A, Valaitis S. Prolapsing vaginal fibroepithelial polyp. BMJ Case Rep 2022; 15(12): e250967. http://dx.doi.org/10.1136/bcr-2022-250967 PMID: 36549754
- [5] Sakai M, Nishimura B, Hiyama T, et al. Imaging of diffuse fibroepithelial polyps on surgical free flap in oral cancer patients: Two case reports. Neuroradiology 2023; 65(4): 815-8. http://dx.doi.org/10.1007/s00234-022-03112-7 PMID: 36622386
- [6] Perez-Milicua G, Orejuela FJ, Patil NM, Adeyemi-Fowode O. Fibroepithelial polyp of the vagina in an adolescent. J Pediatr Adolesc Gynecol 2018; 31(2): 188. http://dx.doi.org/10.1016/j.jpag.2018.02.071
- [7] Cai Y, Zhang Z, Yue X. Rare giant primary ureteral polyp: A case report and literature review. Mol Clin Oncol 2017; 6(3): 327-30. http://dx.doi.org/10.3892/mco.2017.1146 PMID: 28451407
- [8] Zhanghuang C, Hang Y, Ji F, et al. Congenital giant fibroepithelial polyp of the scrotum in an infant: The first case report from China. Front Pediatr 2023; 11: 1191983. http://dx.doi.org/10.3389/fped.2023.1191983 PMID: 37492601
- [9] Park K, Abbas P, Langenburg S, Poulik J, Hanan A, Shehata BM. Giant anal fibroepithelial polyp in a healthy teenage boy: A case report and literature review. Fetal Pediatr Pathol 2022; 41(3): 493-8. http://dx.doi.org/10.1080/15513815.2020.1843575 PMID:

http://dx.doi.org/10.1080/15513815.2020.1843575 PMID 33164595

- [10] Can B, Ozluk AY. Giant fibroepithelial polyps: Why do they grow excessively? Sisli Etfal Hospital 2020; 54(2): 257-60. PMID: 32617069
- [11] Chawla S, Jain S, Kaur L, Gupta B, Rajaram S, Goel N. Giant fibroepithelial polyp: A rare tumour of vulva. IJGO 2017; 15(2): 27. http://dx.doi.org/10.1007/s40944-017-0118-0
- [12] Kurniadi A, Rinaldi A, Yulianti H, Bazar AR, Prasetyawati RD, Tjandraprawira KD. Multiple vulvar giant fibroepithelial polyps: A rare case occurrence. Case Rep Obstet Gynecol 2022; 2022: 1-4. http://dx.doi.org/10.1155/2022/5712925 PMID: 35392425
- [13] Rexhepi M, Trajkovska E, Besimi F, Rufati N. Giant fibroepithelial polyp of vulva: A case report and review of literature. Prilozi 2018; 39(2-3): 127-30. http://dx.doi.org/10.2478/prilozi-2018-0051 PMID: 30864355

[14] Pharaon M, Warrick J, Lynch MC. Fibroepithelial stromal polyp of

the vulva: Case report and review of potential histologic

mimickers. Int J Gynecol Pathol 2018; 37(4): e1-5. http://dx.doi.org/10.1097/PGP.00000000000437 PMID: 28863069

[15] Woo AKM. Depression and anxiety in pain. Rev Pain 2010; 4(1): 8-12.

http://dx.doi.org/10.1177/204946371000400103 PMID: 26527193

- [16] Korkontzelos I, Mpourazanis G, Goshi F, et al. Giant ulcerated fibroepithelial stromal polyp of the vulva: A case report. Cureus 2023; 15(6): e40017. http://dx.doi.org/10.7759/cureus.40017 PMID: 37425539
- [17] Dura MC, Aktürk H, Sungur GŞ, Alsalamin WOI. A giant fibroepithelial polyp of the vulva. Cureus 2023; 15(5): e39152. http://dx.doi.org/10.7759/cureus.39152 PMID: 37378209
- [18] Hernández Martínez AM, Hernández Alarcón R, Fernández Silvestre EC. Vulvar fibroepithelial polyp: A case series. Rev Bras Ginecol Obstet 2023; 69: 1-4.
- [19] Mulik SS, Acharya DB. Large vulval acrochordon in postmenopausal woman—An uncommon presentation: Case report. J South Asian Fed Obstet Gynecol 2023; 15(6): 749-51. http://dx.doi.org/10.5005/jp-journals-10006-2335
- [20] Donata E. Giant fibroepithelial stroma polyp of vulva: A rare case. J Pak Assoc Dermatol 2022; 32: 647-9.
- [21] Kurniawati EM, Djunaidi F, Kurniasari N. Giant fibroepithelial polyps of the vulva in a woman with uterine myoma and primary infertility: A case report and literature review. Am J Case Rep 2022; 23: e933198-1. http://dx.doi.org/10.12659/AJCR.933198 PMID: 35034088
- [22] Smet C, Gomes TG, Silva L, Matias J. Giant fibroepithelial vulvar polyp in a pregnant woman. BMJ Case Rep 2021; 14(1): e236106. http://dx.doi.org/10.1136/bcr-2020-236106 PMID: 33431442
- [23] Quinn M. Large vulvar mass in young adult female. Int J Case Rep Imag 2020; 11: 1. http://dx.doi.org/10.5348/101116Z01MO2020CI
- [24] Yoo J, Je BK, Yeom SK, Park YS, Min KJ, Lee JH. Giant fibroepithelial stromal polyp of the vulva: Diffusion-weighted and conventional magnetic resonance imaging features and pathologic correlation. J Pediatr Adolesc Gynecol 2019; 32(1): 93-7. http://dx.doi.org/10.1016/j.jpag.2018.08.006 PMID: 30149124
- [25] Kumar A, Hasin N, Sinha AK, Kumar S, Bhadani P. Giant fibro epithelial polyp in a young girl: A rare case report. Int J Surg Case Rep 2017; 38: 83-5.
 - http://dx.doi.org/10.1016/j.ijscr.2017.06.059 PMID: 28743098
- [26] Colak E, Ikinci A, Kucuk GO, Kesmer S, Yildirim K. Giant fibroepithelial polyp of the perineum. Int J Surg Case Rep 2015; 17: 126-7.

http://dx.doi.org/10.1016/j.ijscr.2015.11.008 PMID: 26615447

[27] Madueke-Laveaux OS, Gogoi R, Stoner G. Giant fibroepithelial stromal polyp of the vulva: Largest case reported. Ann Surg Innov Res 2013; 7(1): 8.

http://dx.doi.org/10.1186/1750-1164-7-8 PMID: 23842282

- [28] Kesić V, Vieira-Baptista P, Stockdale CK. Early diagnostics of vulvar intraepithelial neoplasia. Cancers 2022; 14(7): 1822. http://dx.doi.org/10.3390/cancers14071822 PMID: 35406594
- [29] Xu C, Zhang Z, Ye H, et al. Imaging diagnosis and endoscopic treatment for ureteral fibroepithelial polyp prolapsing into the bladder. J XRay Sci Technol 2013; 21(3): 393-9. http://dx.doi.org/10.3233/XST-130390 PMID: 24004869
- [30] Heller DS, Day T, Allbritton JI, et al. Diagnostic criteria for differentiated vulvar intraepithelial neoplasia and vulvar aberrant maturation. J Low Genit Tract Dis 2021; 25(1): 57-70. http://dx.doi.org/10.1097/LGT.00000000000572 PMID: 33105449
- [31] Kato H, Kanematsu M, Sato E, Ito N, Furui T, Hirose Y. Magnetic resonance imaging findings of fibroepithelial polyp of the vulva: Radiological-pathological correlation. Jpn J Radiol 2010; 28(8): 609-12.

http://dx.doi.org/10.1007/s11604-010-0465-6 PMID: 20972861

[32] Cicogna S, Dellino M, Miano ST, et al. Aggressive angiomyxoma of the lower female genital tract in pregnancy: A review of the MITO rare tumors group. Cancers 2023; 15(13): 3403.

- [33] Helmi Z, Nori W, Ghani Zghair MA, Abdulrahman Hadi BA. Bartholin gland carcinoma: A case report. J Pak Med Assoc 2021; 71(12): S86-8. PMID: 35130249
- [34] Avila J, Nicol K, Hewitt GD, Matson SC. Vulvar fibroepithelial polyps in a female adolescent: A case report. J Pediatr Adolesc Gynecol 2017; 30(5): 595-7.

http://dx.doi.org/10.1016/j.jpag.2017.04.004 PMID: 28527670

[35] Daneshpajooh A, Mirzaei M, Iranpour M. An unusual presentation of giant fibroepithelial polyp of the vagina. Iran J Pathol 2021; 16(4): 456-60.

http://dx.doi.org/10.30699/ijp.2021.522689.2566 PMID: 34567198

[36] Gangadharan V, M S, S S, P H, P D, Kumari K. Fibroepithelial stromal polyp of vagina in pregnancy - A mimicker. IJCR 2022; 8: 93-5.

http://dx.doi.org/10.32677/ijcr.v8i4.3357

- [37] Wali V, Wali VV. Assessment of various biochemical parameters and BMI in patients with skin tags. J Clin Diagn Res 2016; 10(1): BC09-11.
 PMID: 26894056
- [38] Dane C, Dane B, Cetin A, Erginbas M, Tatar Z. Association of psoriasis and vulval fibroepithelial polyp: First reported case. Am J Clin Dermatol 2008; 9(5): 333-5. http://dx.doi.org/10.2165/00128071-200809050-00008 PMID: 18717609
- [39] El-Sayed ME, Rakha EA, Reed J, Lee AHS, Evans AJ, Ellis IO. Predictive value of needle core biopsy diagnoses of lesions of

uncertain malignant potential (B3) in abnormalities detected by mammographic screening. Histopathology 2008; 53(6): 650-7. http://dx.doi.org/10.1111/j.1365-2559.2008.03158.x PMID: 19076681

- [40] Behtash N, Mousavi A, Tehranian A, Khanafshar N, Hanjani P. Embryonal rhabdomyosarcoma of the uterine cervix: Case report and review of the literature. Gynecol Oncol 2003; 91(2): 452-5. http://dx.doi.org/10.1016/S0090-8258(03)00539-0 PMID: 14599884
- [41] Fife CE, Farrow W, Hebert AA, et al. Skin and wound care in lymphedema patients: A taxonomy, primer, and literature review. Adv Skin Wound Care 2017; 30(7): 305-18. http://dx.doi.org/10.1097/01.ASW.0000520501.23702.82 PMID: 28617750
- [42] Innocent EE-E, Wokpeogu PC, Sudor WB, Nwolim PJ. The effect of ethanolic seed extract of Solanum lycopersicum on wound healing. Eur J Pharm Med Res 2017; 5: 356-61.
- [43] Rogers AH, Farris SG. A meta-analysis of the associations of elements of the fear-avoidance model of chronic pain with negative affect, depression, anxiety, pain-related disability and pain intensity. Eur J Pain 2022; 26(8): 1611-35. http://dx.doi.org/10.1002/ejp.1994 PMID: 35727200
- [44] Lawal BK, Yahya A, Zubair SY, et al. Giant fibro-epithelial polyp of the vulva: A case report. J West Afr Coll Surg 2024; 14(2): 229-32. http://dx.doi.org/10.4103/jwas.jwas 117 23 PMID: 38562392
- [45] Malhotra P, Morgaonkar M, Shetty AR, Rahangdale V. Unveiling the giant: Understanding large vulvar fibroepithelial polyps: A case report with review of literature. NJDVL 2024; 22(2): 35-7.