

The Skin-brain Axis in Cutaneous Aging: Neurobiological Foundations and Clinical Implications



Ahmad Taha Khalaf^{1,*} , Haitham Bader¹ , Waseem H. Q. Alkhaffaf¹ , Yahya D. M. Zakaria¹ , Omar Y. Abdullah¹  and Mohammed N. S. Al-Mallah¹ 

¹Medical College, Nineveh University, Nineveh Governorate 41001, Iraq

Abstract:

The skin-brain axis is a bidirectional neuroimmunoendocrine axis connecting psychological stress, brain signaling, immune responses, and skin homeostasis. A hypothetical link between dysregulation of this network and skin ageing is neurogenic inflammation, oxidative stress, cellular senescence, inflammaging, and activation of the hypothalamic-pituitary-adrenal axis. This dysregulation is increasingly recognized as associated with mechanisms of skin ageing, including neurogenic inflammation, oxidative stress, cellular ageing, inflammaging, and activation of the hypothalamic-pituitary-adrenal axis. This review focuses on the neurobiological mechanisms underlying cutaneous ageing and emphasizes the role of neurocutaneous receptors, neurotransmitters, neuropeptides, and neuro-signaling pathways in the stress response. It also examines emerging trends, such as 'neurocosmetics' including peptides that act on neurocutaneous pathways to stimulate regeneration, stress-modulating peptides, and myorelaxant peptides. This review provides new insight into the skin-brain axis as a potential model for better understanding stress-related skin ageing by combining intrinsic and extrinsic skin ageing with neurobiology. There are a variety of neuroactive cosmetic strategies, with some having mechanistic and preliminary clinical evidence, while others do not and require controlled, randomized studies with objective markers. Future preventive and therapeutic strategies to regulate neuroimmune and neuroendocrine signaling to preserve skin homeostasis and reduce stress-induced skin ageing may be available.

Keywords: Skin-brain axis, Cutaneous aging, Neurogenic inflammation, Neurocosmetics, Oxidative stress, Inflammaging, Neuroendocrine signaling, Cellular senescence.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Medical College, Nineveh University, Nineveh Governorate 41001, Iraq; E-mail: ahmadtaha11@yahoo.com

Cite as: Khalaf A, Bader H, Alkhaffaf W, Zakaria Y, Abdullah O, Al-Mallah M. The Skin-brain Axis in Cutaneous Aging: Neurobiological Foundations and Clinical Implications. Open Dermatol J, 2026; 20: e18743722507190. <http://dx.doi.org/10.2174/0118743722507190261009061047>



CrossMark

Received: July 14, 2026
Revised: September 09, 2026
Accepted: September 16, 2026
Published: October 28, 2026



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

In medical practice, the skin and the nervous system are well known to be connected. However, this connection has only recently begun to shape new concepts in skin physiology. Today, discussion of the "cutaneous nervous system" centers on the skin as an organ of sensory and neuroimmune function that decodes emotional states and environmental signals [1]. This change has laid the foundation for therapeutic and cosmetic strategies that impact the skin's function and psychological state. The key aspect of this emerging field is that keratinocytes,

melanocytes, and resident immune cells can both produce and respond to neurotransmitters, including endorphins, serotonin, dopamine, and substance P. These mediators help regulate barriers, modulate inflammation, and shape sensory perception. This creates a peripheral neuroendocrine network linked to emotional states and stress responses. This role may also help explain why emotional stress can worsen atopic dermatitis, psoriasis, and rosacea, all of which have been associated with disruption of neuroimmune signaling along the skin-brain axis [1, 2]. This relationship is increasingly understood as knowledge of skin microbiomes grows. A disturbance in

microbial balance can activate systemic inflammatory pathways, which can influence Hypothalamic-Pituitary-Adrenal (HPA) activity and consequently affect affective regulation and skin inflammation [1, 3]. Recent studies also suggest that microbial metabolites and antibiotic exposure influence neuro-inflammatory responses that could underlie stress-related skin reactivity [4]. While researchers are still exploring the mechanisms behind these effects, microbiome-targeted topical approaches are gaining traction, with preliminary links established between skin composition and psychological and skin effects [4]. Meanwhile, the notion that topical products could affect emotional health adds complexity. Various agents have been suggested to modulate skin reactivity, increase comfort, and influence subjective sensations about skin condition, such as acetylhexapeptide-8, cannabinoid receptor agonists, and agents related to endorphin signaling [5]. Such statements are at least open to questioning because therapies affecting the neurosensory pathways could also influence emotional reactions. This matters because it is a biochemical effect that is not necessarily what one expects to see physiologically, and a physiologically measurable effect that may reflect a well-documented placebo effect [3-5].

Meanwhile, digital technologies are changing how we define this field. Through innovative artificial intelligence technologies, indicators of emotion can be captured from facial microexpressions, thermopatterns, and circadian or autonomic signals, enabling more personalized and responsive skin care [6]. These technologies can provide benefits but also raise data privacy and governance concerns, as well as other challenges in real-world care delivery. As these are developed, ethical issues need to be considered to ensure that they improve patients' health and do not create new dangers. Taken together, these trends suggest we may be seeing the beginning of a multidisciplinary field spanning dermatology, neuroscience, psychology, sensory biology, and data science. Much of the advancement is theoretical, and far more work is required for these ideas to have clinical impact. Standardized methods, objective parameters, and studies using longer periods are needed in the future, with integration of dermatological and psychophysiological parameters [1, 2]. If this groundwork is laid, emotionally responsive skin care may become a vital aspect of integrative skin care. This field focuses on the dynamic relationship between emotions, the brain, and skin [7]. This convergence is important for the discipline. It broadens the concept of skin as a barrier to an interface inhabited by neurological, immunological, microbial, and psychological processes. Future research on these cutaneous neural pathways, microbially influenced processes, and sensory function could lead to novel approaches to optimizing patient treatments, especially for patients whose skin conditions are closely linked to emotional fluctuations and neuroimmunological sensitivities [8].

Skin ageing is a complex process that involves both genetic ageing mechanisms (intrinsic) and environmental ageing mechanisms (extrinsic) [9]. Although UV exposure and oxidative stress are important, cumulative stress is also important. In recent years, it has been clearly demonstrated that the measurable contribution of psychological stress and interactions between the nervous system and immune system in skin homeostasis are of great importance [9]. This interaction has a developmental basis, and the shared embryologic origin of the epidermis and the nervous system underlies it [10]. Neuroimmunocutaneous models offer mechanistic insight into how emotional and physiological stress affects the aging phenotype [3-5]. Thus, the skin-brain axis has become an exciting field of study again, given its role in modulating skin resilience, inflammatory tone, and regeneration, following the integration of dermatology with neuroscience and immunology.

1.1. Study Design and Literature Search Strategy

This study aimed to conduct a narrative literature review to summarize existing knowledge on the mechanisms in the skin-brain axis involved in cutaneous aging and their possible clinical implications. We searched PubMed/MEDLINE, Scopus, Web of Science, and Google Scholar, focusing on peer-reviewed English-language literature. The following keywords were used in the search: "skin-brain axis," "cutaneous aging," "skin aging," "neurogenic inflammation," "neuroendocrine signaling," "oxidative stress," "inflammaging," "cellular senescence," "neuropeptides," "neurotransmitters," and "neurocosmetics." We included original studies, reviews, and clinical and experimental reports relevant to the mechanisms, biomarkers, and potential therapeutic approaches of the skin-brain axis and aging. We also scanned relevant journals and newspapers to identify additional studies. We synthesized the literature narratively based on the key mechanistic and clinical themes discussed in this review.

2. HISTORY OF THE BRAIN-SKIN LINK

One of the oldest written texts that documents the skin-brain bond is found in the Charaka Samhita of ancient Indian medicine, "Ayurveda," written in the 2nd century CE: the skin (*Twacha*) is closely linked to the mind (*Manas*), as described in the quoted *Chetasamvayi* (inseparable relation) [11]. A similar observation appears in ancient Greek medicine and Avicennian traditions, which recognize a deep, innate bond between the brain and skin, linking them through their shared embryological origin [12]. Hippocrates, in *Epidemics III* (410 BC), reported on the case of Thasos, the wife of Delearches [3]. Following a bout of grief, she had periodic fevers, wrapped herself up, and plucked, scratched, and picked at her hair [13]. Traditional Chinese Medicine views skin disorders as manifestations of internal imbalance. Practitioners use acupuncture and herbal remedies that target the nervous system to restore harmony [14]. In modern medicine, Psychodermatology has emerged as a discipline dedicated to exploring this complex relationship. It aims to

comprehend the bilateral dialogue between the skin and the brain and how treating either alone can lead to incomplete management and early relapses [15].

3. CHARACTERISTICS OF THE SKIN-BRAIN AXIS

3.1. Common Embryonic Origin

The ectoderm contributes to the development of both the skin's epithelial layer and the nervous system [16]. During gastrulation, ectodermal cells, influenced by Wnt, FGF, and BMP signaling pathways, differentiate into either neural or epidermal cells [17]. FGF promotes neural cell formation, while blocking it induces BMP expression and commits ectodermal cells to become epidermal cells [18]. Later, the central neural plate in the ectoderm folds inward to form the neural tube, which develops into the central nervous system. In contrast, the peripheral ectoderm differentiates to form the skin and its appendages. Many molecular cues orchestrate epidermal specification, commitment, stratification, and terminal differentiation.

3.2. Heavy Neural Connection

Because of their shared embryonic development, the brain maintains a strong neural connection with the skin throughout life [19]. The skin is therefore not a simple passive barrier; rather, it is a smart dress for the body and represents its external boundary between the inside and the outside. Therefore, it is regarded as an “external” component of the nervous system [20]. It is estimated that the skin across the whole body is innervated by approximately 230,000 tactile afferents and roughly 200 sensory receptors per cm² [21]. The skin acts as a “peripheral brain” because it has a smart system that can differentiate nociceptive stimuli from pleasant touch using different afferent neurons (unmyelinated (C) and C-tactile fibers, respectively) to the “central brain”. Pain and itching prompt the individual to adopt protective defensive behaviors in response to potential threats [22]. The skin-brain axis is bidirectional, since “non-neural cells modify neural activity” and “sensory neurons influence non-neural cells”. Cytokine release in inflammatory cutaneous disorders induces neuroinflammation and neuroplastic changes, leading to altered mood and cognitive function [23]. Cortisol release is triggered by psychological stress and thus increases inflammatory and pruritic conditions [24, 25]. Beyond skin-mediated mechanisms, stress can also affect the immune system through other pathways, directly impact immune cells, and disrupt barrier function, all of which can worsen different skin disorders [26, 27].

4. MECHANISMS LINKING THE SKIN-BRAIN AXIS TO AGING

4.1. Holistic Theory of Skin Aging

Skin is a Neuro-Immuno-Cutaneous-Endocrine (NICE) system. It has an integrative, bidirectional network between the nervous, immune, and endocrine systems to ensure local and systemic homeostasis. Disequilibrium in this “trinity” function- neuro-, endocrine, and immune-

operates to induce aging [28]. Aging is a multi-faceted, lifelong process involving the interplay between genetically predetermined cellular senescence and various extrinsic age-accelerating stressors [29]. Four pillars drive aging: neurogenic inflammation, oxidative stress, inflammaging, and hypothalamic-pituitary-adrenal axis signaling. Skin aging is a complex process that unfolds throughout the lifespan and results from the interaction between intrinsic cellular aging mechanisms and extrinsic aging factors [30]. Genomic instability results from errors during repeated DNA replication. Telomeres are a biological clock that regulates skin aging. The senescence-Associated Secretory Phenotype (SASP) is a set of pro-inflammatory cytokines secreted by senescent fibroblasts. This leads to excessive reactive oxidative stress and activation of the ATM/p53/p21 signaling pathway, resulting in cell cycle arrest and apoptosis. This process is complex and multifaceted, involving the interplay between innate cellular senescence and multiple extrinsic stressors of ageing [30]. Over time, these factors act together through shared molecular pathways to alter the phenotype and disrupt skin homeostasis. The molecular mechanisms behind these errors include repeated genomic replication, which causes DNA instability [31]. Moreover, Telomere Attrition (shortening of the protective DNA caps at the ends of chromosomes with age) is a biological clock that regulates skin ageing and reduces the chances of developing skin cancer due to DNA damage [31]. Cellular senescence has beneficial protective functions (such as permanent loss of cell division in damaged cells and regulation of tissue remodeling). Still, chronic senescence can be detrimental, including cell-cycle arrest and the establishment of a low-grade, persistent subclinical inflammatory state, “inflamm-aging” [32, 33]. Senescent fibroblasts release excessive amounts of mixed pro-inflammatory cytokines TNF- α , IL-1, IL-6, IFN- γ , collectively known as the “Senescence-Associated Secretory Phenotype (SASP)”, which act in autocrine, paracrine, and endocrine ways [34]. This domino effect forces neighboring healthy cells to become senescent (“zombie cells”) [35] by increasing Reactive Oxidative Stress (ROS) [36]. Oxidative stress will activate the ATM (ataxia telangiectasia mutated)/p53/p21 signaling pathway. This results in cell cycle arrest, senescence, and apoptosis [37]. At the same time, fibroblasts produce excess MMPs that break down collagen and the extracellular matrix [38]. Sustained exposure to harsh environmental factors gradually dysregulates barrier function and the repair system, leading to premature skin aging. Oxidative stress is thought to be the primary cause of extrinsic skin aging, in which a mismatch between abundant intracellular free-radical production and limited antioxidant defenses occurs. Consequently, unstable chemical ROS are overproduced [39]. ROS damage mitochondrial membranes, reduce ATP synthesis, and trigger a cascade of events that accelerate skin aging. In addition, it increases Matrix Metalloproteinase (MMP) expression, which degrades collagen and other extracellular matrix components. Thus, inhibiting MMPs is key to maintaining youthful skin [40]. Senescent

fibroblasts reduce the production of Insulin-Like Growth Factor 1 (IGF-1), a hormone that signals keratinocytes to repair from UV-induced DNA damage. The internal glycosylation process contributes to protein structural damage and to the skin-aging phenotype *via* a “Maillard-like reaction”. Excess reduced sugars and free amino groups of proteins, DNA, and lipids bind nonenzymatically to form harmful Advanced Glycation End products (AGEs). This process forms an irreversibly cross-linked, stiff protein that appears wrinkled, yellow, and dull. AGE accumulation also accelerates skin aging by increasing ROS generation and activating matrix-degrading enzymes (metalloproteinases), which fragment collagen and the matrix. Additionally, skin acts as a stress sensor, modulating homeostasis *via* the neuroendocrine and immune systems and signaling molecules [41-43]. It protects against stressors through keratinocyte differentiation, the pigmentary system, and communication with the dermis/hypodermis. Dysfunctions can cause various skin disorders and aging.

4.2. Inflamm-aging

Lifetime exposure to harsh environmental factors makes it difficult for the skin to repair itself and cope with stressors. Dyshomeostasis of barrier functions will induce autoreactivity, immunosenescence, and premature aging [44]. At the same time, repeated genomic mitosis leads to error accumulation, DNA instability, and telomere attrition [40-43]. The biological clock leads to the irreversible cessation of division in old cells (senescence). Senescent cells do not die; instead, they remain metabolically active and release harmful substances [45]. This maladaptive state induces subclinical persistent inflammation, “inflamm-aging” [46]. Senescent fibroblasts release a Senescence-Associated Secretory Phenotype (SASP) of pro-inflammatory cytokines [47]. SASP includes TNF- α , IL-1 β , IL-6, and IFN- γ , which act in autocrine, paracrine, and endocrine ways. Senescent cells behave like “zombie cells,” causing nearby healthy cells to become senescent [48]. This domino effect is attributed to excessive Reactive Oxygen Species (ROS) production. Oxidative stress activates the ATM (Ataxia Telangiectasia Mutated)/p53/p21 signaling pathway [49]. This results in cell cycle arrest, senescence, and apoptosis. At the same time, fibroblasts produce excess matrix metalloproteinases (MMPs) that break down collagen and the extracellular matrix [50].

4.3. Neurogenic Inflammation

The interaction of extrinsic environmental hazards and intrinsic physical or psychological stresses causes “stress-induced aging of the skin” [51]. Dysregulated neurotrophin release and pro-inflammatory loops drive aging. Mast cell activation and degranulation play a “key role” in neurogenic inflammation [52]. They are the “initiators and amplifiers” of inflammation through bidirectional signaling with peripheral neurons [53]. Stress induces the release of neuropeptides responsible

for itching and scratching, such as Calcitonin Gene-Related Peptide (CGRP) and substance P (SP). The neurotransmitters activate mast cells directly through the mast-related G protein-coupled receptor X2 (MRGPRX2) or indirectly through the neurokinin-1 receptor (NK-1R) [54]. Activated MCs undergo degranulation, releasing potent mixtures of tryptase, chymase, cytokines, chemokines, and histamine. Tryptase binds to protease-activated receptor-2 (PAR-2) on nerve endings, leading to the release of additional neuropeptides [55]. Patients enter a vicious circle that intensifies neurogenic inflammation and pruritus by positively fueling the mast cell-nerve loop (SP induces MC degranulation, and MC mediators enhance SP release) [56].

4.4. Oxidative Stress

Oxidative stress is thought to be the primary cause of extrinsic skin aging. It results from an imbalance between excessive intracellular free radical production and limited antioxidant defenses [57]. Consequently, Reactive Oxygen Species (ROS) become unstable and are overproduced. ROS accelerate skin aging by damaging the mitochondrial membrane, reducing ATP synthesis, and triggering an inflammatory cascade [58]. In addition, they increase MMP expression, which degrades collagen and other Extracellular Matrix (ECM) components. Thus, inhibiting MMPs is key to maintaining youthful skin [59]. Increased ECM breakdown and remodeling will minimize skin resilience, exacerbate inflammation, and disrupt the cutaneous immune cell orchestra [57, 58]. As fibroblasts age, they stop producing insulin-like growth factor 1 (IGF-1), which signals keratinocytes to repair UV-damaged DNA [60]. Stress signaling increases ROS generation in keratinocytes and fibroblasts. Oxidative stress causes DNA damage, telomere loss, and reduced DNA repair capacity. Neuroimmune imbalance could therefore work alongside photoaging. Changes in the skin's microbial flora may affect the cytokine network and nerve sensitization. The mechanisms remain unclear, but new evidence indicates that microbial metabolites can affect inflammatory signaling and stress responses, which can influence aging outcomes. The major interconnected mechanisms linking psychological and environmental stress to cutaneous aging through neuroendocrine, inflammatory, oxidative, and cellular senescence pathways are illustrated in Fig. (1).

Extrinsic age-accelerating stressors and genetically programmed cellular senescence affect the Neuro-Immuno-Cutaneous-Endocrine (NICE) system through neurogenic inflammation and oxidative stress, the concept of “inflammaging,” and Hypothalamic-Pituitary-Adrenal (HPA) axis signaling. These interacting pathways lead to increased release of neuropeptides, mast cell activation, production of Reactive Oxygen Species (ROS), cellular senescence, secretion of the Senescence-Associated Secretory Phenotype (SASP), release of matrix metalloproteinases (MMPs), and degradation of the Extracellular Matrix (ECM), all of which collectively contribute to the skin-aging phenotype.




Fig. (1). Neuro-immuno-cutaneous interactions in stress-related skin aging.

4.5. Hypothalamic-pituitary-adrenal Signaling

Psychological stress activates the Hypothalamic-Pituitary-Adrenal (HPA) axis, a key player in the skin-brain dialogue, leading to the systemic release of glucocorticoids, mainly cortisol. This is adaptive in the short term, but long-term activation burdens cutaneous structure and function. High glucocorticoid levels have been shown to delay barrier recovery after disruption, primarily by interfering with lipid synthesis and lamellar body secretion from the stratum corneum [61]. At the cellular level, cortisol inhibits keratinocyte proliferation and differentiation by promoting epidermal thinning and decreased turnover, and it inhibits fibroblast activity and collagen production in the dermis [62]. This is further enhanced by reduced transforming growth factor- β signaling and increased matrix metalloproteinase expression, which favor extracellular matrix degradation [62]. Importantly, the skin itself contains a peripheral HPA axis (with the ability to locally produce corticotropin-releasing hormone, adrenocorticotrophic hormone, and even cortisol, thereby amplifying the stress signal within the cutaneous microenvironment) [63]. These local NE networks indicate that stress-associated aging is not just a systemic process but also a tissue-specific one. Both central and peripheral HPA signaling are chronically

dysregulated and are associated with characteristics of premature skin aging, such as decreased elasticity, impaired wound healing, and a higher incidence of inflammatory dermatoses. These findings suggest that HPA axis activity is an important mechanistic connection between psychological stress and structural and functional decline in aging skin, and that therapies that restore neuroendocrine balance may have therapeutic potential.

5. NEUROCUTANEOUS RECEPTORS

The skin harbors a wide variety of distinct receptor types involved in barrier function, immune surveillance, and tissue homeostasis, which guide the direction of aging. While classical receptor systems generally amplify inflammatory signaling and oxidative stress, transient receptor potential channels have a greater influence in this regard, and cannabinoid receptors have a counter-regulatory effect on homeostasis. Serotonin and dopamine receptors also seem to promote cellular resilience, thereby enhancing keratinocyte viability and wound healing [64]. In addition, so-called ectopic receptors in the skin may participate in other physiological functions in regeneration, such as promoting keratinocyte migration and fibroblast activity, as well as modulating inflammation and local tissue tone through acetylcholine receptors. At

the translational level, any approach that modulates TRP signaling pathways to reduce activity and/or augment or mimic cannabinoid and/or monoaminergic signaling pathways is a candidate for restoring a more youthful cellular environment. Similarly, selective modulation of ectopic receptors can restore the skin to a reparative and/or quiescent state, thus highlighting their potential role in new skin rejuvenation strategies. The details of these receptors are shown below:

5.1. Transient Receptor Potential (TRP) Channels

These are cutaneous, multimodal, Ca²⁺-permeable sensory ion channels that act as a neuro-inflammatory hub [65]. They consist of 6 families: TRPA, TRPC, TRPM, TRPML, TRPP, and TRPV. They respond to temperature alterations, UV radiation, and pH changes. Overactivation of TRP induces chronic inflammation and accelerated skin aging [66]. TRP is widely expressed in nerve endings, keratinocytes, melanocytes, and immune cells. It acts as a sensory "Switch" at the nerve ending: a UV sensor, a skin-sensitivity inducer, and an age accelerator. Irritant activation of TRP channels at nerve endings leads to SP release, which exacerbates neurogenic inflammation, induces MMPs, and accelerates skin aging [67]. Neurocosmetics are designed to desensitize TRP channels, reduce inflammation, and restore the skin barrier.

5.2. Cannabinoid Receptors (CB)

Cannabinoid receptors are part of the group of G Protein-Coupled Receptors (GPCRs) [68]. CBs are important in maintaining skin homeostasis, regulating the immune system, and reducing inflammation [69]. The cannabinoid receptors produce endogenous "anti-aging" effects by dampening inflammation, countering free radicals, and improving cell regeneration [70]. Insufficiency decreases the skin's antioxidant activity, disrupts mitochondrial function, increases pro-inflammatory markers, promotes skin aging, thins the skin, and impairs wound healing [71]. This suggests topical products containing cannabinoids are potential anti-aging agents, without the psychotropic properties of oral or inhaled cannabis [72].

5.3. Serotonergic and Dopaminergic Receptors in Skin Aging

The skin is a neurohormonal organ that receives and processes neurotransmitters (such as serotonin and dopamine). These neurotransmitters are crucial but often overlooked in skin aging. Serotonin Receptors, such as 5 HT, are expressed in keratinocytes, fibroblasts, and melanocytes. It acts as a natural antioxidant, aids wound healing, reduces inflammation, and pigmentation [73]. Dopamine and its two receptor types (D1 and D2) signaling maintain skin vitality by regulating cell proliferation, sebum production, and immune response [74]. Dopamine behaves differently in wound healing depending on the receptor type involved. D1 receptor signaling enhances angiogenesis and wound healing, while D2 receptor activation hinders them [75]. Serotonin and dopamine, along with their receptors, decline

proportionally with age [76]. This will reduce their protective, anti-inflammatory, and metabolic-regulating effects. "Skin is a mirror of the soul." Chronic stress depletes these neurotransmitters, accelerating skin aging [77]. Imbalances in serotonin-dopamine pathways are linked to inflammation and pigment disorders. Moreover, manipulating these pathways (such as using serotonin and dopamine antagonists at specific doses) may mimic the effects of calorie restriction on individuals' longevity.

5.4. Olfactory Receptors (ORs)

It is the largest G protein-coupled chemosensory receptor family ectopically expressed in extranasal tissues, including skin cells (melanocytes, fibroblasts, hair follicle epithelial cells, and keratinocytes) [78]. These receptors play a crucial role in various skin functions, from enhancing skin health to accelerating wound healing [79]. OR receptors are highly expressed and promote basal cell proliferation, a process essential for wound healing. It prevents skin aging by suppressing oxidative stress in keratinocytes [80]. It also plays an essential role in fibroblast survival and collagen synthesis [81]. In addition to restoring skin barrier function [82], experimental studies show that Sandalore enhances keratinocyte function in wound healing, whereas the antagonist oxyphenylon inhibits this effect [83].

5.5. Acetylcholine Receptors

Acetylcholine (ACh) receptors are members of the superfamily of neurotransmitter-gated ion channels. They are expressed in two forms: nicotinic (nAChR) and muscarinic (mAChR). Their natural ligand is the endogenous neurotransmitter acetylcholine (ACh). They classically exist in neural tissues. Recently, extraneuronal localization has been recognized in many tissues, including skin cells [84-86]. Moreover, cells that transiently reside in the skin, such as immune system cells, are involved. Exploration of these receptors is a promising target for managing skin inflammation and aging [84-86]. In neural tissues, ACh released from the nerve terminal fuses with the cell membrane *via* a SNARE complex. Released ACh binds to nAChR to induce muscle contraction. Wrinkles can be reduced by relaxing facial muscle contractions. This can be achieved by inhibiting SNARE complex formation or by direct binding to nAChRs. The non-neuronal cholinergic system regulates Epidermal Homeostasis, glandular function, and immune response. nAChR agonists suppress UV-induced proinflammatory cytokine expression and oxidative stress, both of which are critically involved in dermal photoaging.

6. NEUROTRANSMITTERS AND NEUROACTIVE AGENTS

Skin aging is not just a structural derangement but a complex neurobiological process. Many neurotransmitters and bioactive molecules mediate this process and can activate, inhibit, or modulate it. Imbalances dysregulate the skin microenvironment and contribute to early skin aging. There are four groups of neurotransmitters: Monoamines (Serotonin, Dopamine, Norepinephrine,

Epinephrine, Histamine), Amino acids (GABA, Glutamate, Glycine), Neuropeptides (Substance P, Opioids, Neuropeptide Y), and others (Acetylcholine). They may be excitatory, inhibitory, or modulatory inputs to effectors. The table summarizes the cutaneous neurotransmitters important for skin homeostasis and aging processes [84-86].

The skin is now recognized as a functional neuroendocrine system that can synthesize and respond to neurotransmitters independently of central input. Keratinocytes, melanocytes, and dermal fibroblasts express the enzymatic machinery required for local monoamine production, including serotonin and catecholamines, and environmental stressors such as ultraviolet radiation and oxidative stress can dynamically modulate their levels [87, 88]. This local signaling network appears to fine-tune key processes relevant to aging, including keratinocyte proliferation, melanogenesis, and microvascular regulation. In parallel, acetylcholine, produced within the non-neuronal cholinergic system, acts as a widespread regulator of epidermal differentiation, barrier formation, and immune responses, further supporting the concept that skin neurotransmitters function as local cytoremitters rather than solely as neural messengers [89]. Such observations suggest that alterations in cutaneous neurotransmitter balance may precede visible structural changes, positioning them as early drivers of age-related dysfunction rather than secondary phenomena.

At the tissue level, neuroactive mediators converge on pathways that define the aging phenotype, particularly chronic low-grade inflammation, extracellular matrix degradation, and impaired repair capacity. Ultraviolet exposure induces matrix metalloproteinases, notably MMP-1, which initiate collagen breakdown and disrupt dermal architecture, a process further amplified by senescent fibroblasts through the release of pro-inflammatory and proteolytic factors [90, 91]. Within this context, neuropeptides such as substance P exhibit dual, context-dependent effects, promoting tissue repair and angiogenesis under physiological conditions but showing diminished efficacy with advancing age [92, 93].

Conversely, histamine has been shown to impair keratinocyte differentiation and weaken barrier integrity, linking neurogenic inflammation to functional decline of the epidermis [94, 95]. Emerging data also point to a protective role for inhibitory neurotransmitters; GABAergic signaling in dermal fibroblasts has been associated with enhanced extracellular matrix synthesis and improved resistance to oxidative stress [96]. Taken together, these findings reinforce the notion that neurotransmitter imbalance contributes directly to the molecular and cellular events underlying cutaneous aging. However, further work is needed to translate these insights into targeted therapeutic strategies. The principal mediators of the Neuro-Immuno-Cutaneous-Endocrine (NICE) network, their physiological roles, and their potential contributions to skin aging are summarized in Table 1.

7. NEUROCOSMETICS AND THE EXPANDING BIOLOGICAL HORIZON OF COSMETIC SCIENCE

The emergence of neurocosmetics marks a conceptual departure in skincare, moving from purely aesthetic, surface-level enhancement of skin as a “tissue” to a holistic, systematic approach that treats skin as a “neuroimmunologic organ” [97]. While traditional cosmetics primarily provide symptomatic management that addresses physical appearance (*e.g.*, epidermal hydration, barrier repair, and visible texture), neurocosmetics interact directly with the skin-brain axis by modulating neurotransmitter release and receptor signaling, thereby addressing stress response and skin homeostasis. As the comparative framework summarizes, this paradigm shift moves beyond structural enhancement toward systemic modulation, aiming to reinforce skin resilience, reduce stress-induced flares, and potentially influence psychocutaneous interactions. The transition reflects a more integrated biological understanding of aging and reactivity, positioning neurocosmetics within a holistic model that bridges dermatologic physiology and neurobiology rather than confining intervention to the epidermal surface alone. Table 2 summarizes the differences between traditional cosmetics and neurocosmetics.

Table 1. The role and dysregulation of key mediators of the neuro-immuno-cutaneous-endocrine (NICE) network in skin aging.

Mediators	Primary Source	Physiological Roles	Impact on Skin Aging
Substance P / CGRP	Sensory nerve ending	Mast cell degranulation and immune reaction	Neurogenic inflammation, collagen & ECM breaks & remodeling
Serotonin (5-HT)	Systemic and local in keratinocytes	Regulates cell survival and melanogenesis	Skin cell apoptosis, impaired barrier, oxidative stress, and uneven pigmentation.
Dopamine	Local and Sympathetic nerve	Regulates pigmentation, stress, and vasoconstriction.	Increases cellular senescence, Irregular melanin synthesis, and barrier disruption.
Acetyl choline	Neuron and keratinocytes	Muscle contraction, skin adhesion and differentiation	High activity: dynamic expression lines; low activity: delays barrier repair
GABA	Fibroblast	Accelerates cutaneous barrier recovery	Alters muscle tone, shifts inflammation, and decreases collagen, elastin, and hydration.
β -endorphin	Keratinocytes, Appendages.	Regulate differentiation, migration, and barrier	Impaired regeneration, reduced barrier function, itch and skin aging

Table 2. The difference between traditional cosmetics and neurocosmetics.

Features	Traditional Cosmetics	Neurocosmetics
Primary goal	Symptomatic treatment of aging signs	Systematic modulation of the mechanism of aging
Core mechanism	Improve structural integrity	Modulate neurotransmitters and receptor signaling
Biological focus	Surface-oriented visible changes	Deep behavioral alteration of the sensory neuron
Addressed issue	Treat symptom	Treat system
Typical ingredients	Humectant emollients	Neurotransmitter mimickers
Scope of impact	Primary physical and aesthetics	Holistic therapy
Primary goal	Enhance external appearance	Boost skin behavior, resilience and mood
Overstimulation	Irritation	Calm and resilient skin

8. ANTIAGING NEUROCOSMETICS

Neurocosmetics are an advanced, non-toxic, bioactive skincare category designed to modify skin-brain axis neurons and receptors to mitigate skin aging [98]. These products utilize short-chain biomimetic ingredients that act locally to relax muscles, soothe, or regenerate skin. The first group has “Botox-like” effects that target the neuromuscular junction by inhibiting ACh and reducing facial muscle expression. The second group ameliorates chronic inflammation, collagen breakdown, and skin-barrier impairment induced by local HPA-axis activation by reducing local cortisol levels (a danger signal) and increasing endorphin release (feel-good neuropeptides). The last group improves regeneration by restarting cellular repair through enhanced cellular crosstalk within the skin-brain axis, thereby enhancing collagen, elastin, and ECM synthesis.

8.1. Myo-relaxant to Relax Expression Lines

Wrinkles are the consequence of lifelong muscle contraction. The process begins with acetylcholine release at the neuromuscular junction. SNARE complexes dock

ACh-containing vesicles to facilitate their fusion with the neuronal membrane. Released ACh binds to postsynaptic nAChRs, initiating muscle contraction. Several neurotransmitter inhibitor peptides have been designed with sequences that resemble and compete with these key steps (SNAP-25, nAChR, and ion channels). These Botox-like myorelaxant peptides mitigate wrinkle depth in a non-invasive, effective, and safe manner. They satisfy growing consumer demand among injection-averse individuals, younger individuals focusing on prejuvenation, and as a post-Botox treatment to extend treatment results. Although the efficacy of topical Botox-like peptides varies widely based on formulation, concentration, and skin barrier penetration, they can be ranked in descending order as follows: Syn-Ake (Tripeptide-3), Vialox (Pentapeptide-3), XEP-018 (or XEP-030/XEP-018), Snap-8 (Acetyl Octapeptide-3), Argireline (Acetyl Hexapeptide-8), Myoxinol (Hibiscus Esculentus Extract), and Leuphasyl (Pentapeptide-18). Table 3 summarizes the proven Botox-like myorelaxant peptides, in descending order of efficacy, for reducing wrinkle depth after 4 weeks of continuous daily use.

Table 3. Neurocosmetics act as myorelaxants, ranked in descending order according to the percentage of wrinkle depth reduction after continuous daily use for 4 weeks.

Name	Ingredient	Source	Year	Developer	Mechanism of Action	Wrinkle Depth Reduction	Ref
Syn-Ake	Tripeptide	Snake venom mimic	2006	DSM	nAChR antagonist and modulates GABAA receptors	52%	[99]
Vialox	Pentapeptide-3	Synthetic	2003	DSM	Cura-like antagonist of nAChRs	49%	[100]
XEP-30 and XEP-018	μ -conotoxin CnIIIC McMahon	marine cone snail venom mimic	2014	Activen	Blocks ACh release by targeting Voltage-gated sodium channels	48%	[101]
Snap-8	Acetyl Octapeptide-3 Ji	Synthetic extract	2003	Lipotec	Elongated Argireline SNARE destabilizer	38%	[102]
Argireline	Acetyl Hexapeptide-8	Synthetic extract	2001	Lipotec	SNARE destabilizer and blocks ACh release	30%	[103]
Myoxinol	okra extract	Natural extract	2002	BASF	Enhancing GABAergic transmission	26%	[104]
Leuphasyl	Pentapeptide-18	Mimic enkephalin	2005	Lipotec	blocking calcium channels	24%	[105]

8.2. Neuro-relaxant to Combat Stress-inducing Skin Aging

Chronic stress accelerates skin aging by activating the local HPA axis and the 11 β -HSD1 enzyme, which converts inactive cortisone to biologically active cortisol, a “stress hormone,” leading to increased glycation and structural damage of collagen and elastin, catabolic degradation of dermal proteins, and a reduction in the natural moisturizer hyaluronic acid, thereby impairing the skin barrier. The body counteracts this by releasing β -endorphins, a “skin wellness neurotransmitter,” from the degradation of POMC. The Complex β -endorphin/ μ -opiate inhibits signal transmission from nociceptors, fine-tunes the immune response, accelerates skin regeneration, and promotes wound healing. Neuro-relaxing anti-aging neurocosmetics ingredients address the “stress-aging” cycle *via* the skin-brain axis by targeting the local HPA axis, inhibiting the 11 β -HSD1 enzyme, or boosting “feel-good” neurotransmitters, such as endorphins, to improve skin

appearance. Table 4 summarizes the proven neuro-relaxant neurocosmetics available commercially.

8.3. Skin Regeneration Boosters

Regenerative skin boosters and neurocosmetics target a biological decline that accelerates after age 50: increased epidermal turnover, fibroblast senescence, declining collagen and elastin levels, and ECM damage. Neurocosmetics rejuvenation boosters are formulated with specialized peptides that help restore skin structure, resilience, and barrier function. They contain signal peptides (which stimulate fibroblasts to increase collagen, elastin, fibronectin, and hyaluronic acid through the TGF- β pathway), carrier peptides (which transport the metal ions copper and manganese, which are involved in wound healing, collagen synthesis, and antioxidant defense), and enzyme-inhibitor peptides (which inhibit MMPs to prevent collagen degradation). Representative regenerative peptides and neurocosmetic ingredients, together with their proposed mechanisms of action, are summarized in Table 5.

Table 4. Neuro-relaxants to reduce stress-induced skin aging.

Name	Source	Developer	Year	Action	Ref
Agascalm	<i>Agastache mexicana</i>	Provital	2015	Prevents NF- κ B release; decreases redness and TEWL; reduces cytokine release	[106]
Betaphroline	<i>Tephrosia purpurea</i> (wild indigo)	Givaudan	2012	Stimulates β -endorphin, Pain relief, and improvement in well-being	[107]
Happybelle-PE	<i>Vitex agnus-castus</i> extract	Mibelle	2005	Increases collagen I and III; improves skin hydration and firmness; decreases wrinkle depth.	[108]
Happy Skin	<i>Rhodiola rosea</i> root extract	Provital	2013	Protection from free-radical-induced damage; Increase in endorphin levels	[109]
Neurophroline	seeds of <i>Tephrosia purpurea</i> extract	Givaudan	2016	Inhibition of cortisol release; stimulation of β -endorphin production	[110]
Sepicalm S WP	<i>Nymphaea alba</i> flower extract	Seppic	2006	β -endorphin production increases skin water content and decreases redness.	[111]
Nirvanesque	<i>Paeonia suffruticosa</i> extract	NUXE	2022	Produces enkephalins as a “neuro-relaxing” molecule	[112]
NeuroCalm™	<i>Peonia suffruticosa</i>	LANCÔME	2026	Soothing day cream moisturizes and nourishes	[113]
Defensil SOFT	Edible mushroom <i>Albatrellus ovinus</i>	Rahn	2017	Blocks TRPV1 receptor; reduces skin hyperalgesia and neurogenic inflammation	[114]
Mariliance	<i>Rhodorus marinus</i> , a red microalga	Givaudan	2015	Reduce TRPV1 expression, release of IL-1 α , & sensitivity	[114]
Neutrazen	synthetic tripeptide	Lucas Meyer Cosmetics	2000	Biomimetic of (POMC) reduces SP, IL-1, IL-8, and neurogenic inflammation.	[115]
Ocaline PF	<i>Cucurbita pepo</i> Seed Extract	Givaudan	2025	Preventive effect on sensitive skin/stinging, soothing of irritated skin	[116]
Pinolumin	Swiss stone pine wood	Mibelle	2016	Inhibit release of (CGRP, IL-8, and PGE2), act on TRPV1; improvement skin tone	[117]
SensAmon P5	Synthetic pentapeptide from sea anemone	Mibelle	2017	Inhibition of TRPV1 receptor activation; reduction of skin sensitivity.	[115]
Sensityl	<i>Phaeodactylum tricornutum</i>	Givaudan	2019	skin soothing; control of the acute phase of inflammation, immune cell recruitment and inflammatory cytokines	[118]
Skinasensyl LS 9749	Synthetic Acetyl Tetrapeptide-15	BASF	2008	Binding to the μ -opioid receptor; inhibition of CGRP release; increased skin tolerance threshold.	[115]
Sepicalm S WP	water lily extract	Seppic	2006	production of β -endorphins; inhibition hyaluronidase; Increase skin water, reduce redness and stinging sensation	[119]

Table 5. Neurocosmetics that act as skin boosters to regenerate aged skin.

Name	Ingredient	Year	Developer	Mechanism of Action	Ref
Matrixyl®	Palmitoyl Pentapeptide-4	2000	Sederma	Stimulates collagen and ECM components	[120]
Decorinyl™	Tripeptide-10 Citrulline	2010	Lipotec SAU	Regulates fibrillogenesis	[121]
Biopeptide El™	Palmitoyl Hexapeptide-12	1992	Sederma	Promotes collagen, elastin and glycosaminoglycans production.	[122]
Syn®-Coll	Palmitoyl Tripeptide-5	2012	dsm-firmenich	Boosts collagen production by stimulating TGF-β production	[123]
Dermican™	Acetyl Tetrapeptide 9	2011	BASF	Stimulates collagen type I and lumican synthesis.	[123]
Syniorage™	Acetyl Tetrapeptide-11	2013	BASF	Accelerates keratinocyte growth and syndecan-1 synthesis.	[123]
Lipospondin	Elaidoyl Tripeptide-37	2010	Sederma	Stimulates TGF-β, promoting collagen and elastin production.	[123]
ND	Peptamide-6 /Hexapeptide-11	2012	Lonza Personal Care	Modulates key genes responsible for collagen synthesis, improving skin elasticity.	[123]
Progeline™	Trifluoroacetyl Tripeptide-2	2012	Lucas Meyer Cosmetics	Inhibits progerin synthesis, a protein that accelerates cell aging.	[123]
Rigin™	Palmitoyl Tetrapeptide-7	2011	Sederma	Decreases IL-6, stimulates laminins IV and V and collagen VII	[123]
Perfection Peptide P7™	Heptapeptide	2014	Mibelle Biochemistry	Stimulating Nrf2-dependent antioxidant enzymes.	[123]
GEKG	Tetrapeptide-21	2011	Evonik Industries	Stimulates collagen synthesis at the mRNA and protein levels	[123]
KP1	SA 1-III	2020	Università di Firenze	Modulates collagen turnover.	[123]
No	Soybean peptides	2015	Shenzhen	An enzyme-inhibitor peptide neutralizes free radicals, increases collagen production, and offers UV protection.	[123]
PeptAlde™ 4.0	Rice peptides	2020	BASF	Enzyme-inhibiting peptide, Antioxidant, and anti-inflammatory effects.	[123]
Activated Silk™ peptides	Silk peptides	2020s	Nature	An enzyme inhibitor peptide inhibits MMP-1 and stimulates hyaluronan synthase 2 gene expression.	[123]
Cu-GHK	Copper Tripeptide-1	1990s	Dr. Loren Pickart.	Carrier-regenerative peptide improves healing, modulates MMP-1, promotes collagen and glycosaminoglycans, and has antioxidant properties.	[123]
GHK-Mn2+	Manganese Tripeptide-1	1980s	Dr. Loren Pickart.	Carrier-regenerative peptide improves healing, acts as an antioxidant, and reduces hyperpigmentation.	[123]

9. LIMITATIONS AND KNOWLEDGE GAPS

Other regeneration-based strategies, such as stem cell-derived extracellular vesicles and growth factor-based formulations, could also indirectly stabilize the neuroimmune axis by reducing inflammation and boosting dermal repair. The next step in research should be to assess whether these interventions can alter stress-related signaling networks in senescent skin. The field is conceptually sound, but it has limitations. Multiple studies are in development, but they include small sample sizes or surrogate markers. It is difficult to differentiate expectancy-induced placebo effects from biochemical modulation. Additionally, stressors vary by person, leading to different microbiome compositions; each person's neuroendocrine system also responds differently to stressors, making generalization difficult. Standardized protocols, reproducible biomarkers, and longitudinal designs are required to determine whether modulating the skin-brain axis has long-term benefits.

CONCLUSION

The skin-brain axis is an important framework for conceptualizing the interplay between psychological stress, neuroimmune signaling, oxidative stress, and inflammation in skin ageing. The new generation of neuroactive/regenerative therapies has promise for maintaining skin homeostasis and counteracting stress-induced ageing effects. However, evidence remains limited and inconsistent. More well-designed clinical trials

are needed to determine the therapeutic significance of targeting the skin-brain interaction in ageing skin using objective biomarkers.

AUTHORS' CONTRIBUTIONS

The authors confirm their contributions to the paper as follows: A.T.K. and H.B.: Study conception and design; W.H.Q.A., Y.D.M.Z., and O.Y.A.: Literature collection and data acquisition; A.T.K., H.B., and M.N.S.A.M.: Analysis and interpretation of the literature and findings; A.T.K., W.H.Q.A.: Drafting of the manuscript; A.T.K., H.B., Y.D.M.Z., O.Y.A., M.N.S.A.M.: Critical revision of the manuscript. All authors reviewed the manuscript, approved the final version, and agreed to be accountable for the work.

LIST OF ABBREVIATIONS

MMP = Matrix Metalloproteinase

ROS = Reactive Oxidative Stress

TRP = Transient Receptor Potential

HPA = Hypothalamic-Pituitary-Adrenal

NICE = Neuro-Immuno-Cutaneous-Endocrine

SASP = Senescence-Associated Secretory Phenotype

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

Dr. Ahmad Taha Khalaf is the Editorial Advisory Board member of The Open Dermatology Journal.

ACKNOWLEDGEMENTS

The authors thank Shamil Mahmood Ali Al-Jobore, Al-Salam Teaching Hospital, Mosul, Nineveh Governorate, Iraq, for his valuable support during the revision of the manuscript.

REFERENCES

- [1] Vidal Yucha SE, Tamamoto KA, Kaplan DL. The importance of the neuro-immuno-cutaneous system on human skin equivalent design. *Cell Prolif* 2019; 52(6): 12677. <http://dx.doi.org/10.1111/cpr.12677> PMID: 31441145
- [2] Scholzen T, Armstrong CA, Bunnett NW, Luger TA, Olerud JE, Ansel JC. Neuropeptides in the skin: Interactions between the neuroendocrine and the skin immune systems. *Exp Dermatol* 1998; 7(2-3): 81-96. <http://dx.doi.org/10.1111/j.1600-0625.1998.tb00307.x> PMID: 9583747
- [3] Slominski A, Wortsman J. Neuroendocrinology of the Skin. *Endocr Rev* 2000; 21(5): 457-87. <http://dx.doi.org/10.1210/edrv.21.5.0410> PMID: 11041445
- [4] Guo Z, Yang J, Zang R, Yang Y, Wang Q, Xu C. The brain-gut-skin axis in inflammatory and disfiguring skin diseases: Mechanistic insights, clinical correlations, and therapeutic strategies. *Front Immunol* 2026; 17: 1737303. <http://dx.doi.org/10.3389/fimmu.2026.1737303> PMID: 41836406
- [5] Joachim RA, Kuhlmei A, Dinh QT, *et al.* Neuronal plasticity of the "brain-skin connection": Stress-triggered up-regulation of neuropeptides in dorsal root ganglia and skin *via* nerve growth factor-dependent pathways. *J Mol Med* 2007; 85(12): 1369-78. <http://dx.doi.org/10.1007/s00109-007-0236-8> PMID: 17639286
- [6] Kumar G, Das T, Singh K. Early detection of depression through facial expression recognition and electroencephalogram-based artificial intelligence-assisted graphical user interface. *Neural Comput Appl* 2024; 36(12): 6937-54. <http://dx.doi.org/10.1007/s00521-024-09437-z>
- [7] Christopoulos GI, Uy MA, Yap WJ. The body and the brain: Measuring skin conductance responses to understand the emotional experience. *Organ Res Methods* 2019; 22(1): 394-420. <http://dx.doi.org/10.1177/1094428116681073>
- [8] O'Sullivan RL, Lipper G, Lerner EA. The neuro-immuno-cutaneous-endocrine network: Relationship of mind and skin. *Arch Dermatol* 1998; 134(11): 1431-5. <http://dx.doi.org/10.1001/archderm.134.11.1431> PMID: 9828880
- [9] Hussein RS, Bin Dayel S, Abahusseini O, El-Sherbiny AA. Influences on skin and intrinsic aging: Biological, environmental, and therapeutic insights. *J Cosmet Dermatol* 2025; 24(2): 16688. <http://dx.doi.org/10.1111/jocd.16688> PMID: 39604792
- [10] Sarnat HB, Flores-Sarnat L. Embryology of the neural crest: Its inductive role in the neurocutaneous syndromes. *J Child Neurol* 2005; 20(8): 637-43. <http://dx.doi.org/10.1177/08830738050200080101> PMID: 16225807
- [11] Pandey MM, Rastogi S, Rawat AKS. Indian traditional ayurvedic system of medicine and nutritional supplementation. *Evid Based Complement Alternat Med* 2013; 2013: 1-12. <http://dx.doi.org/10.1155/2013/376327> PMID: 23864888
- [12] Ghezloo S, Daneshfard B, Khadem E. Early description of the common embryonic origin of the skin and nervous system by Avicenna. *Galen Med J* 2024; 13: 3330. <http://dx.doi.org/10.31661/gmj.v13i.3330> PMID: 39224546
- [13] Waas R, Yesudian P. Plucking, picking, and pulling: The hair-raising history of trichotillomania. *Int J Trichology* 2018; 10(6): 289-90. http://dx.doi.org/10.4103/ijt.ijt_69_18 PMID: 30783339
- [14] He T, Zheng Y, Yan J, *et al.* Skin-Brain Axis: Neural pathways in acupuncture treatment. *Chin Med* 2025; 20(1): 163. <http://dx.doi.org/10.1186/s13020-025-01213-y> PMID: 41053779
- [15] Jafferany M, Franca K. Psychodermatology: Basics concepts. *Acta Derm Venereol* 2014; 96(217): 0. <http://dx.doi.org/10.2340/00015555-2378> PMID: 27282585
- [16] Jameson C, Boulton KA, Silove N, Nanan R, Guastella AJ. Ectodermal origins of the skin-brain axis: A novel model for the developing brain, inflammation, and neurodevelopmental conditions. *Mol Psychiatry* 2023; 28(1): 108-17. <http://dx.doi.org/10.1038/s41380-022-01829-8> PMID: 36284159
- [17] Itoh K, Sokol S. Early development of epidermis and neural tissue. In: Moody SA, Ed. *Principles of developmental genetics*. (2nd ed.). Academic Press 2015; pp. 189-201. <http://dx.doi.org/10.1016/B978-0-12-405945-0.00011-9>
- [18] Zhu XJ, Liu Y, Dai ZM, *et al.* BMP-FGF signaling axis mediates Wnt-induced epidermal stratification in developing mammalian skin. *PLoS Genet* 2014; 10(10): 1004687. <http://dx.doi.org/10.1371/journal.pgen.1004687> PMID: 25329657
- [19] Talagas M. Anatomical contacts between sensory neurons and epidermal cells: An unrecognized anatomical network for neuro-immuno-cutaneous crosstalk. *Br J Dermatol* 2023; 188(2): 176-85. <http://dx.doi.org/10.1093/bjd/ljac066> PMID: 36763869
- [20] Rabischong P. The skin neural interface. In: Ajeena IM, Ed. *Advances in clinical neurophysiology*. IntechOpen 2012. <http://dx.doi.org/10.5772/51601>
- [21] Roosterman D, Goerge T, Schneider SW, Bunnett NW, Steinhoff M. Neuronal control of skin function: The skin as a neuroimmunoendocrine organ. *Physiol Rev* 2006; 86(4): 1309-79. <http://dx.doi.org/10.1152/physrev.00026.2005> PMID: 17015491
- [22] Tan CC, Soh KV, Wang E, Choi ECE. The brain-skin connection: A narrative review of neuroendocrine and immune pathways. *JAAD Int* 2026; 24: 112-23. <http://dx.doi.org/10.1016/j.jdin.2025.10.008> PMID: 41393358
- [23] Ferreira BR, Aguirre CC, Rapoport-Hubschman N, *et al.* The skin-brain connection and pleasant touch as supportive care for psychocutaneous disorders. *Skin Health Dis* 2023; 4(1): 310. PMID: 38312257
- [24] Yoon KN, Chung JH. Healthy skin, Healthy brain. *J Dermatol Sci* 2025; 119(2): 47-52. <http://dx.doi.org/10.1016/j.jdermsci.2025.06.001> PMID: 40681402
- [25] Steinhoff M, Dréno B, Rey AB, Akimichi M. The skin-brain dialogue: Advancing psychodermatology through integrated approaches. *J EADV Clin Pract* 2025; 4(S1): S49-58. <http://dx.doi.org/10.1002/jvc2.70100>
- [26] Chen Y, Lyga J. Brain-skin connection: Stress, inflammation and skin aging. *Inflamm Allergy Drug Targets* 2014; 13(3): 177-90. <http://dx.doi.org/10.2174/1871528113666140522104422> PMID: 24853682
- [27] Lyu F, Wu T, Bian Y, Zhu K, Xu J, Li F. Stress and its impairment of skin barrier function. *Int J Dermatol* 2023; 62(5): 621-30. <http://dx.doi.org/10.1111/ijd.16598> PMID: 36759891
- [28] Jin R, Luo L, Zheng J. The trinity of skin: Skin homeostasis as a neuro-endocrine-immune organ. *Life* 2022; 12(5): 725. <http://dx.doi.org/10.3390/life12050725> PMID: 35629392
- [29] Liang Y, Su W, Wang F. Skin ageing: A progressive, multi-factorial condition demanding an integrated, multilayer-targeted remedy. *Clin Cosmet Investig Dermatol* 2023; 16: 1215-29. <http://dx.doi.org/10.2147/CCID.S408765> PMID: 37192990
- [30] Wong QYA, Chew FT. Defining skin aging and its risk factors: A systematic review and meta-analysis. *Sci Rep* 2021; 11(1): 22075. <http://dx.doi.org/10.1038/s41598-021-01573-z> PMID: 34764376
- [31] Potapova T, Gorbysky G. The consequences of chromosome segregation errors in mitosis and meiosis. *Biology* 2017; 6(1): 12. <http://dx.doi.org/10.3390/biology6010012> PMID: 28208750

- [32] Liu Z, Sun C, Zhang Z, Jiang Y, Zhao C. Telomeres in skin aging. *Biogerontology* 2025; 26(2): 83.
<http://dx.doi.org/10.1007/s10522-025-10228-9> PMID: 40159528
- [33] Huang W, Hickson LJ, Eirin A, Kirkland JL, Lerman LO. Cellular senescence: The good, the bad and the unknown. *Nat Rev Nephrol* 2022; 18(10): 611-27.
<http://dx.doi.org/10.1038/s41581-022-00601-z> PMID: 35922662
- [34] Pilkington SM, Bulfone-Paus S, Griffiths CEM, Watson REB. Inflammaging and the Skin. *J Invest Dermatol* 2021; 141(4) (Suppl. 1): 1087-95.
<http://dx.doi.org/10.1016/j.jid.2020.11.006> PMID: 33358020
- [35] Lopes-Paciencia S, Saint-Germain E, Rowell MC, Ruiz AF, Kalegari P, Ferbeyre G. The senescence-associated secretory phenotype and its regulation. *Cytokine* 2019; 117: 15-22.
<http://dx.doi.org/10.1016/j.cyt.2019.01.013> PMID: 30776684
- [36] Rius-Pérez S, Torres-Cuevas I, Millán I, Ortega ÁL, Pérez S. PGC-1 α , inflammation, and oxidative stress: An integrative view in metabolism. *Oxid Med Cell Longev* 2020; 2020: 1-20.
<http://dx.doi.org/10.1155/2020/1452696> PMID: 32215168
- [37] Guo Z, Kozlov S, Lavin MF, Person MD, Paull TT. ATM activation by oxidative stress. *Science* 2010; 330(6003): 517-21.
<http://dx.doi.org/10.1126/science.1192912> PMID: 20966255
- [38] Papaccio F, D Arino A, Caputo S, Bellei B. Focus on the contribution of oxidative stress in skin aging. *Antioxidants* 2022; 11(6): 1121.
<http://dx.doi.org/10.3390/antiox11061121> PMID: 35740018
- [39] Feng C, Chen X, Yin X, Jiang Y, Zhao C. Matrix metalloproteinases on skin photoaging. *J Cosmet Dermatol* 2024; 23(12): 3847-62.
<http://dx.doi.org/10.1111/jocd.16558> PMID: 39230065
- [40] Wandery F, Schmid D. Senolytics: Eliminating “zombie cells” in the skin. A novel anti-aging mechanism to combat senescent cells. *HPC Today* 2020; 15(5): 18-20.
- [41] Chen C, Zhang JQ, Li L, *et al.* Advanced glycation end products in the skin: Molecular mechanisms, methods of measurement, and inhibitory pathways. *Front Med* 2022; 9: 837222.
<http://dx.doi.org/10.3389/fmed.2022.837222> PMID: 35646963
- [42] Peng H, Gao Y, Zeng C, *et al.* Effects of Maillard reaction and its product AGEs on aging and age-related diseases. *Food Sci Hum Wellness* 2024; 13(3): 1118-34.
<http://dx.doi.org/10.26599/FSHW.2022.9250094>
- [43] Stammers M, Ivanova IM, Niewczas IS, *et al.* Age-related changes in the physical properties, cross-linking, and glycation of collagen from mouse tail tendon. *J Biol Chem* 2020; 295(31): 10562-71.
<http://dx.doi.org/10.1074/jbc.RA119.011031> PMID: 32381510
- [44] Chambers ES, Vukmanovic-Stejic M. Skin barrier immunity and ageing. *Immunology* 2020; 160(2): 116-25.
<http://dx.doi.org/10.1111/imm.13152> PMID: 31709535
- [45] Regulski MJ. Cellular senescence: What, why, and how. *Wounds* 2017; 29(6): 168-74.
PMID: 28682291
- [46] Di Micco R, Krizhanovsky V, Baker D, d'Adda di Fagagna F. Cellular senescence in ageing: From mechanisms to therapeutic opportunities. *Nat Rev Mol Cell Biol* 2021; 22(2): 75-95.
<http://dx.doi.org/10.1038/s41580-020-00314-w> PMID: 33328614
- [47] Pajak J, Nowicka D, Szepletowski JC. Inflammaging and immunosenescence as part of skin aging—A narrative review. *Int J Mol Sci* 2023; 24(9): 7784.
<http://dx.doi.org/10.3390/ijms24097784> PMID: 37175491
- [48] Kumari R, Jat P. Mechanisms of cellular senescence: Cell cycle arrest and senescence-associated secretory phenotype. *Front Cell Dev Biol* 2021; 9: 645593.
<http://dx.doi.org/10.3389/fcell.2021.645593> PMID: 33855023
- [49] Xie X, Zhang Y, Wang Z, *et al.* ATM at the crossroads of reactive oxygen species and autophagy. *Int J Biol Sci* 2021; 17(12): 3080-90.
<http://dx.doi.org/10.7150/ijbs.63963> PMID: 34421351
- [50] Quan T, Little E, Quan H, Qin Z, Voorhees JJ, Fisher GJ. Elevated matrix metalloproteinases and collagen fragmentation in photodamaged human skin: Impact of altered extracellular matrix microenvironment on dermal fibroblast function. *J Invest Dermatol* 2013; 133(5): 1362-6.
<http://dx.doi.org/10.1038/jid.2012.509> PMID: 23466932
- [51] Parrado C, Mercado-Saenz S, Perez-Davo A, Gilaberte Y, Gonzalez S, Juarranz A. Environmental stressors on skin aging: Mechanistic insights. *Front Pharmacol* 2019; 10: 759.
<http://dx.doi.org/10.3389/fphar.2019.00759> PMID: 31354480
- [52] Polsky LR, Rentscher KE, Carroll JE. Stress-induced biological aging: A review and guide for research priorities. *Brain Behav Immun* 2022; 104: 97-109.
<http://dx.doi.org/10.1016/j.bbi.2022.05.016> PMID: 35661679
- [53] Naguib Abdel Hafez SM, Ali ESMN, Naguib SM. The role of mast cells in accelerating our skin aging. *J Clin Images Med Case Rep* 2021; 2(5): 1345.
<http://dx.doi.org/10.52768/2766-7820/1345>
- [54] Marek-Jozefowicz L, Nedoszytko B, Grochocka M, *et al.* Molecular mechanisms of neurogenic inflammation of the skin. *Int J Mol Sci* 2023; 24(5): 5001.
<http://dx.doi.org/10.3390/ijms24055001> PMID: 36902434
- [55] Hsin L, Fernandopulle NA, Ding J, *et al.* The effect of substance P and its common *in vivo*-formed metabolites on MRGPRX2 and human mast cell activation. *Pharmacol Res Perspect* 2022; 10(4): 00990.
<http://dx.doi.org/10.1002/prp2.990> PMID: 35904495
- [56] Xu K, Wang L, Lin M, He G. Update on protease-activated receptor 2 in inflammatory and autoimmune dermatological diseases. *Front Immunol* 2024; 15: 1449126.
<http://dx.doi.org/10.3389/fimmu.2024.1449126> PMID: 39364397
- [57] Gupta K, Harvima IT. Mast cell-neural interactions contribute to pain and itch. *Immunol Rev* 2018; 282(1): 168-87.
<http://dx.doi.org/10.1111/imr.12622> PMID: 29431216
- [58] Rinnerthaler M, Bischof J, Streubel M, Trost A, Richter K. Oxidative stress in aging human skin. *Biomolecules* 2015; 5(2): 545-89.
<http://dx.doi.org/10.3390/biom5020545> PMID: 25906193
- [59] Liguori I, Russo G, Curcio F, *et al.* Oxidative stress, aging, and diseases. *Clin Interv Aging* 2018; 13: 757-72.
<http://dx.doi.org/10.2147/CIA.S158513> PMID: 29731617
- [60] Kemp M, Spandau D, Travers J. Impact of age and insulin-like growth factor-1 on DNA damage responses in UV-irradiated human skin. *Molecules* 2017; 22(3): 356.
<http://dx.doi.org/10.3390/molecules22030356> PMID: 28245638
- [61] Franceschi C, Garagnani P, Parini P, Giuliani C, Santoro A. Inflammaging: A new immune-metabolic viewpoint for age-related diseases. *Nat Rev Endocrinol* 2018; 14(10): 576-90.
<http://dx.doi.org/10.1038/s41574-018-0059-4> PMID: 30046148
- [62] Maarouf M, Maarouf CL, Yosipovitch G, Shi VY. The impact of stress on epidermal barrier function: An evidence-based review. *Br J Dermatol* 2019; 181(6): 1129-37.
<http://dx.doi.org/10.1111/bjd.17605> PMID: 30614527
- [63] Vukelic S, Stojadinovic O, Pastar I, *et al.* Cortisol synthesis in epidermis is induced by IL-1 and tissue injury. *J Biol Chem* 2011; 286(12): 10265-75.
<http://dx.doi.org/10.1074/jbc.M110.188268> PMID: 21239489
- [64] Pondeljak N, Lugović-Mihic L. Stress-induced interaction of skin immune cells, hormones, and neurotransmitters. *Clin Ther* 2020; 42(5): 757-70.
<http://dx.doi.org/10.1016/j.clinthera.2020.03.008> PMID: 32276734
- [65] Gupta D, Kaushik D, Mohan V. Role of neurotransmitters in the regulation of cutaneous wound healing. *Exp Brain Res* 2022; 240(6): 1649-59.
<http://dx.doi.org/10.1007/s00221-022-06372-0> PMID: 35488904
- [66] Gouin O, L'Herondelle K, Lebonvallet N, *et al.* TRPV1 and TRPA1 in cutaneous neurogenic and chronic inflammation: Pro-inflammatory response induced by their activation and their sensitization. *Protein Cell* 2017; 8(9): 644-61.
<http://dx.doi.org/10.1007/s13238-017-0395-5> PMID: 28364279
- [67] Kang SM, Han S, Oh JH, *et al.* A synthetic peptide blocking TRPV1 activation inhibits UV-induced skin responses. *J Dermatol Sci* 2017; 88(1): 126-33.
<http://dx.doi.org/10.1016/j.jdermsci.2017.05.009> PMID: 28551094

- [68] Xiao T, Sun M, Zhao C, Kang J. TRPV1: A promising therapeutic target for skin aging and inflammatory skin diseases. *Front Pharmacol* 2023; 14: 1037925. <http://dx.doi.org/10.3389/fphar.2023.1037925> PMID: 36874007
- [69] Yoo EH, Lee JH. Cannabinoids and their receptors in skin diseases. *Int J Mol Sci* 2023; 24(22): 16523. <http://dx.doi.org/10.3390/ijms242216523> PMID: 38003712
- [70] Tortolani D, Di Meo C, Standoli S, *et al.* Rare phytocannabinoids exert anti-inflammatory effects on human keratinocytes via the endocannabinoid system and MAPK signaling pathway. *Int J Mol Sci* 2023; 24(3): 2721. <http://dx.doi.org/10.3390/ijms24032721> PMID: 36769042
- [71] Liu C, Li H, Xu F, Jiang X, Ma H, Seeram NP. Cannabidiol protects human skin keratinocytes from hydrogen peroxide-induced oxidative stress via modulation of the caspase-1-IL-1 β axis. *J Nat Prod* 2021; 84(5): 1563-72. <http://dx.doi.org/10.1021/acs.jnatprod.1c00083> PMID: 33955754
- [72] Leal EC, Moura LIF, Pirzgalska RM, *et al.* Diabetes and Cannabinoid CB1 receptor deficiency promote similar early onset aging-like changes in the skin. *Exp Gerontol* 2021; 154: 111528. <http://dx.doi.org/10.1016/j.exger.2021.111528> PMID: 34437952
- [73] Redmond MC, Finn DP. The therapeutic potential of cannabidiol in skin conditions. *J Cosmet Dermatol* 2025; 24(11): 70527. <http://dx.doi.org/10.1111/jocd.70527> PMID: 41178006
- [74] Enkhtaivan E, Lee CH. Role of amine neurotransmitters and their receptors in skin pigmentation: Therapeutic implication. *Int J Mol Sci* 2021; 22(15): 8071. <http://dx.doi.org/10.3390/ijms22158071> PMID: 34360837
- [75] Tammaro A, Cavallotti C, Gaspari AA, Narcisi A, Parisella FR, Cavallotti C. Dopaminergic receptors in the human skin. *J Biol Regul Homeost Agents* 2012; 26(4): 789-95. PMID: 23241131
- [76] Vaughn A, Davis M, Sivamani R, Isseroff R. A concise review of the conflicting roles of dopamine-1 *versus* dopamine-2 receptors in wound healing. *Molecules* 2017; 23(1): 50. <http://dx.doi.org/10.3390/molecules23010050> PMID: 29278360
- [77] Yin JA, Liu XJ, Yuan J, Jiang J, Cai SQ. Longevity manipulations differentially affect serotonin/dopamine level and behavioral deterioration in aging *Caenorhabditis elegans*. *J Neurosci* 2014; 34(11): 3947-58. <http://dx.doi.org/10.1523/JNEUROSCI.4013-13.2014> PMID: 24623772
- [78] Yegorov YE, Poznyak AV, Nikiforov NG, Sobenin IA, Orekhov AN. The link between chronic stress and accelerated aging. *Biomedicine* 2020; 8(7): 198. <http://dx.doi.org/10.3390/biomedicine8070198> PMID: 32645916
- [79] Geng R, Wang Y, Fang J, *et al.* Ectopic odorant receptors responding to flavor compounds in skin health and disease: Current insights and future perspectives. *Crit Rev Food Sci Nutr* 2022; 63(28): 9392-408. <http://dx.doi.org/10.1080/10408398.2022.2064812> PMID: 35445618
- [80] Seo J, Choi S, Kim H, Park SH, Lee J. Association between olfactory receptors and skin physiology. *Ann Dermatol* 2022; 34(2): 87-94. <http://dx.doi.org/10.5021/ad.2022.34.2.87> PMID: 35450315
- [81] Kim JS, Lee HL, Jeong JH, *et al.* OR2AT4, an ectopic olfactory receptor, suppresses oxidative stress-induced senescence in human keratinocytes. *Antioxidants* 2022; 11(11): 2180. <http://dx.doi.org/10.3390/antiox11112180> PMID: 36358552
- [82] Son B, Kang W, Park S, Choi D, Park T. Dermal olfactory receptor OR51B5 is essential for survival and collagen synthesis in human dermal fibroblast (Hs68 cells). *Int J Mol Sci* 2021; 22(17): 9273. <http://dx.doi.org/10.3390/ijms22179273> PMID: 34502185
- [83] Kang W, Son B, Park S, Choi D, Park T. UV-irradiation- and inflammation-induced skin barrier dysfunction is associated with the expression of olfactory receptor genes in human keratinocytes. *Int J Mol Sci* 2021; 22(6): 2799. <http://dx.doi.org/10.3390/ijms22062799> PMID: 33802009
- [84] Busse D, Kudella P, Grüning NM, *et al.* A synthetic sandalwood odorant induces wound-healing processes in human keratinocytes via the olfactory receptor OR2AT4. *J Invest Dermatol* 2014; 134(11): 2823-32. <http://dx.doi.org/10.1038/jid.2014.273> PMID: 24999593
- [85] Kurzen H, Berger H, Jäger C, *et al.* Phenotypical and molecular profiling of the extraneuronal cholinergic system of the skin. *J Invest Dermatol* 2004; 123(5): 937-49. <http://dx.doi.org/10.1111/j.0022-202X.2004.23425.x> PMID: 15482483
- [86] Stegemann A, Böhm M. Targeting the $\alpha 7$ nicotinic acetylcholine receptor-A novel road towards the future treatment of skin diseases. *Exp Dermatol* 2020; 29(9): 924-31. <http://dx.doi.org/10.1111/exd.14173> PMID: 32780438
- [87] Slominski AT, Kim TK, Kleszczynski K, *et al.* Characterization of serotonin and *N*-acetylserotonin systems in the human epidermis and skin cells. *J Pineal Res* 2020; 68(2): 12626. <http://dx.doi.org/10.1111/jpi.12626> PMID: 31770455
- [88] Slominski AT, Slominski RM, Raman C, Chen JY, Athar M, Elmets C. Neuroendocrine signaling in the skin with a special focus on the epidermal neuropeptides. *Am J Physiol Cell Physiol* 2022; 323(6): C1757-76. <http://dx.doi.org/10.1152/ajpcell.00147.2022> PMID: 36317800
- [89] Slominski A, Wortsman J, Tobin DJ. The cutaneous serotonergic/melatonergic system: Securing a place under the sun. *FASEB J* 2005; 19(2): 176-94. <http://dx.doi.org/10.1096/fj.04-2079rev> PMID: 15677341
- [90] Zheng J, Wang S, Sun J, Lv J. Dermal fibroblast senescence: The central hub of skin aging—From intrinsic dysfunction to microenvironmental remodeling. *Int J Mol Sci* 2026; 27(4): 1653. <http://dx.doi.org/10.3390/ijms27041653> PMID: 41751788
- [91] Lai CF, Esam G, Lin YC, Chung RJ. From destruction to protection: Rethinking MMP-1 in skin aging and anti-aging strategies. *Arch Dermatol Res* 2026; 318(1): 56. <http://dx.doi.org/10.1007/s00403-025-04518-y>
- [92] Kurzen H, Wessler I, Kirkpatrick C, Kawashima K, Grando S. The non-neuronal cholinergic system of human skin. *Horm Metab Res* 2007; 39(2): 125-35. <http://dx.doi.org/10.1055/s-2007-961816> PMID: 17326008
- [93] Quan T, Qin Z, Xia W, Shao Y, Voorhees JJ, Fisher GJ. Matrix-degrading metalloproteinases in photoaging. *J Invest Dermatol Symp Proc* 2009; 14(1): 20-4. <http://dx.doi.org/10.1038/jidsymp.2009.8> PMID: 19675548
- [94] Zaarour RF, Saha D, Dey R, *et al.* The neuropeptide Substance P facilitates the transition from an inflammatory to proliferation phase-associated responses in dermal fibroblasts. *Exp Dermatol* 2022; 31(8): exd.14573. <http://dx.doi.org/10.1111/exd.14573> PMID: 35353932
- [95] Gschwandtner M, Mildner M, Mlitz V, *et al.* Histamine suppresses epidermal keratinocyte differentiation and impairs skin barrier function in a human skin model. *Allergy* 2013; 68(1): 37-47. <http://dx.doi.org/10.1111/all.12051> PMID: 23157658
- [96] Ito K, Tanaka K, Nishibe Y, Hasegawa J, Ueno H. GABA-synthesizing enzyme, GAD67, from dermal fibroblasts: Evidence for a new skin function. *Biochim Biophys Acta, Gen Subj* 2007; 1770(2): 291-6. <http://dx.doi.org/10.1016/j.bbagen.2006.09.017> PMID: 17113713
- [97] Nan L, Guo P, Hui W, Xia F, Yi C. Recent advances in dermal fibroblast senescence and skin aging: Unraveling mechanisms and pioneering therapeutic strategies. *Front Pharmacol* 2025; 16: 1592596. <http://dx.doi.org/10.3389/fphar.2025.1592596> PMID: 40606604
- [98] Jadhav VM, Kadu VB. Neurocosmetics: Bridging neurosciences and skincare. *Res J Top Cosmet Sci* 2025; 16(2): 124-4. <http://dx.doi.org/10.52711/2321-5844.2025.00020>
- [99] Gok B, Budama-Kilinc Y, Kecel-Gunduz S. Anti-aging activity of Syn-Ake peptide by *in silico* approaches and *in vitro* tests. *J Biomol Struct Dyn* 2024; 42(10): 5015-29. <http://dx.doi.org/10.1080/07391102.2023.2223681> PMID: 37349941
- [100] Nguyen TTM, Yi EJ, Jin X, *et al.* Sustainable dynamic wrinkle

- efficacy: Non-invasive peptides as the future of Botox alternatives. *Cosmetics* 2024; 11(4): 118.
<http://dx.doi.org/10.3390/cosmetics11040118>
- [101] del Río-Sancho S, Cros C, Coutaz B, Cuendet M, Kalia YN. Cutaneous iontophoresis of μ -conotoxin CnIIIC—A potent Na_v 1.4 antagonist with analgesic, anaesthetic and myorelaxant properties. *Int J Pharm* 2017; 518(1-2): 59-65.
<http://dx.doi.org/10.1016/j.ijpharm.2016.12.054> PMID: 28034736
- [102] Ji M, Lee HS, Kim Y, et al. Method development for acetyl octapeptide-3 analysis by liquid chromatography-tandem mass spectrometry. *J Anal Sci Technol* 2020; 11(1): 34.
<http://dx.doi.org/10.1186/s40543-020-00232-8>
- [103] Grosicki M, Latacz G, Szopa A, Cukier A, Kieć-Kononowicz K. The study of cellular cytotoxicity of argireline - An anti-aging peptide. *Acta Biochim Pol* 2014; 61(1): 29-32.
<http://dx.doi.org/10.18388/abp.2014.1919> PMID: 24644551
- [104] Tan C, Lee M, Lim Z, et al. 1172 Targeting skin photoaging with okra seed extracts. *J Invest Dermatol* 2023; 143(5) (Suppl.): S201.
<http://dx.doi.org/10.1016/j.jid.2023.03.1185>
- [105] Dragomirescu A, Andoni M, Ionescu D, Andrei F. The efficiency and safety of Leuphasyl—A Botox-like peptide. *Cosmetics* 2014; 1(2): 75-81.
<http://dx.doi.org/10.3390/cosmetics1020075>
- [106] Nechita MA, Toiu A, Benedec D, et al. *Agastache* species: A comprehensive review on phytochemical composition and therapeutic properties. *Plants* 2023; 12(16): 2937.
<http://dx.doi.org/10.3390/plants12162937> PMID: 37631149
- [107] De Tollenaere M, Meunier M, Scandolera A, et al. Well-aging: A new strategy for skin homeostasis under multi-stressed conditions. *J Cosmet Dermatol* 2020; 19(2): 444-55.
<http://dx.doi.org/10.1111/jocd.13047> PMID: 31232507
- [108] Puglia LT, Lowry J, Tamagno G. *Vitex agnus castus* effects on hyperprolactinaemia. *Front Endocrinol* 2023; 14: 1269781.
<http://dx.doi.org/10.3389/fendo.2023.1269781> PMID: 38075075
- [109] Li Y, Pham V, Bui M, et al. *Rhodiola rosea* L.: an herb with anti-stress, anti-aging, and immunostimulating properties for cancer chemoprevention. *Curr Pharmacol Rep* 2017; 3(6): 384-95.
<http://dx.doi.org/10.1007/s40495-017-0106-1> PMID: 30393593
- [110] Palbag S, Dey BK, Singh NK. Ethnopharmacology, phytochemistry and pharmacology of *Tephrosia purpurea*. *Chin J Nat Med* 2014; 12(1): 1-7.
[http://dx.doi.org/10.1016/S1875-5364\(14\)60001-7](http://dx.doi.org/10.1016/S1875-5364(14)60001-7) PMID: 24484589
- [111] Liao Z, Huo Y, Tan H, Duan F. Cosmetic applications of *Nymphaea* spp.: A review of extraction techniques, functional components, and efficacy mechanisms in skincare. *J Dermatol Sci Cosmet Technol* 2026; 3(1): 100152.
<http://dx.doi.org/10.1016/j.jdsct.2026.100152>
- [112] Li Y, Han J, Gong R, et al. Protective effects of *Paeonia suffruticosa* callus extract in skin through anti-inflammation and repair UVB -induced damage. *Int J Cosmet Sci* 2025; 47(4): 652-64.
<http://dx.doi.org/10.1111/ics.13055> PMID: 40079176
- [113] Liu J, Liu Y, Wang F, Yuan Y, Ma H, Qu L. *Paeonia suffruticosa* Andrews root extract ameliorates photoaging via regulating IRS1/PI3K/FOXO pathway. *Front Pharmacol* 2025; 16: 1520392.
<http://dx.doi.org/10.3389/fphar.2025.1520392> PMID: 40115263
- [114] Scandolera A, Hubert J, Humeau A, et al. GABA and GABA-alanine from the red microalgae *Rhodorus marinus* exhibit a significant neuro-soothing activity through inhibition of neuro-inflammation mediators and positive regulation of TRPV1-related skin sensitization. *Mar Drugs* 2018; 16(3): 96.
<http://dx.doi.org/10.3390/md16030096> PMID: 29562624
- [115] Resende DISP, Ferreira MS, Sousa-Lobo JM, Sousa E, Almeida IF. Usage of synthetic peptides in cosmetics for sensitive skin. *Pharmaceuticals* 2021; 14(8): 702.
<http://dx.doi.org/10.3390/ph14080702> PMID: 34451799
- [116] Grasu AE, Senn R, Halbsguth C, Schenk A, Butterweck V, Miron A. Profiling hydrophilic cucurbita pepo seed extracts: A study of european cultivar variability. *Plants* 2025; 14(15): 2308.
<http://dx.doi.org/10.3390/plants14152308> PMID: 40805657
- [117] Gugerli F, Brodbeck S, Bebi P, et al. Swiss stone pine - Portrait of a mountain forest tree. WSL: Fact sheet 2023.
<http://dx.doi.org/10.55419/wsl:32467>
- [118] Russo MT, Rogato A, Jaubert M, Karas BJ, Falcatore A. *Phaeodactylum tricornutum* : An established model species for diatom molecular research and an emerging chassis for algal synthetic biology. *J Phycol* 2023; 59(6): 1114-22.
<http://dx.doi.org/10.1111/jpy.13400> PMID: 37975560
- [119] Dulal MA, Jahan I, Rasul MG, Islam MR, Khan M, Shah AKMA. Antioxidant and antimicrobial potential of water lily extracts and their effects on the quality of frozen Nile tilapia (*Oreochromis niloticus*) filets. *Food Sci Nutr* 2023; 11(1): 535-44.
<http://dx.doi.org/10.1002/fsn3.3084> PMID: 36655067
- [120] Choi YL, Park EJ, Kim E, Na DH, Shin YH. Dermal stability and *in vitro* skin permeation of collagen pentapeptides (KTTKS and palmitoyl-KTTKS). *Biomol Ther* 2014; 22(4): 321-7.
<http://dx.doi.org/10.4062/biomolther.2014.053> PMID: 25143811
- [121] Raikou V, Varvaresou A, Panderi I, Papageorgiou E. The efficacy study of the combination of tripeptide-10-citrulline and acetyl hexapeptide-3. A prospective, randomized controlled study. *J Cosmet Dermatol* 2017; 16(2): 271-8.
<http://dx.doi.org/10.1111/jocd.12314> PMID: 28150423
- [122] Ligorio C, Tavasoli E, Karaman-Jurukovska N, et al. Noninvasive monitoring of palmitoyl hexapeptide-12 in human skin layers: Mechanical interaction with skin components and its potential skincare benefits. *ACS Appl Bio Mater* 2025; 8(3): 2340-55.
<http://dx.doi.org/10.1021/acsabm.4c01816> PMID: 39964201
- [123] Pintea A, Manea A, Pintea C, et al. Peptides: Emerging candidates for the prevention and treatment of skin senescence: A review. *Biomolecules* 2025; 15(1): 88.
<http://dx.doi.org/10.3390/biom15010088> PMID: 39858482

DISCLAIMER: Please note that this article is currently in the 'Early view' stage and is not the final 'Version of Record'. While it has been accepted, copy-edited, and formatted, however, it is still undergoing proofreading and corrections by the authors. Therefore, the text may still change before the final publication. Although 'Early view' may not have all bibliographic details available, the DOI and the year of online publication can still be used to cite them. The article title, DOI, publication year, and author(s) should all be included in the citation format. Once the 'Final Version of Record' becomes available the 'Early view' will be replaced by that.