



The Open Dermatology Journal

Content list available at: <https://opendermatologyjournal.com>

Supplementary Material



Efficacy of Supramolecular Salicylic Acid 30% Combined with Intense Pulsed Light Compare to Monotherapy Supramolecular Salicylic Acid 30% in Acne Vulgaris Patients

Asoka Ariyawati^{1,*}, Yangmei Chen¹, Yihuan Pu¹, Yujie Zhang¹, Lingzhao Zhang¹, Xinyi Shao¹, Jiayi Chen¹ and Jin Chen¹

¹Department of Dermatology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China




Fig. (S1). Porphyrins of patient no. (23) in the IPL + SSA30% group. (A) Pre-treatment; (B) Post-treatment 4 week; (C) Post-treatment 8 week. Showing improvement after 2 sessions from 53.13% to 74.30%.




Fig. (S2). Red area of patient no. (23) in the IPL + SSA30% group. (A) Pre-treatment; (B) Post-treatment 4 week; (C) Post-treatment 8 week. Showing improvement after 2 sessions from 26.29% to 39.09%.




Fig. (S3). Porphyrins of patient no. (20) in the SSA30% group. (A) Pre-treatment; (B) Post-treatment 4 week; (C) Post-treatment 8 week. Showing improvement after 4 sessions from 56.78% to 72.47%.




Fig. (S4). Red area of patient no. (20) in the SSA30% group. (A) Pre-treatment; (B) Post-treatment 4 week; (C) Post-treatment 8 week. Showing improvement after 4 sessions from 24.12% to 31.73%.

© 2023 Ariyawati *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.